

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

P. ENT COOPERATION TREA

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
 US Department of Commerce
 United States Patent and Trademark
 Office, PCT
 2011 South Clark Place Room
 CP2/5C24
 Arlington, VA 22202
 ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 20 June 2001 (20.06.01)	
International application No. PCT/US00/12811	Applicant's or agent's file reference PF-0693 PCT
International filing date (day/month/year) 10 May 2000 (10.05.00)	Priority date (day/month/year) 11 May 1999 (11.05.99)
Applicant BANDMAN, Olga et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
 28 November 2000 (28.11.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

2r

The International Bureau of WIPO
 34, chemin des Colombettes
 1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

F. Baechler


Telephone No.: (41-22) 338.83.38

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



14

Applicant's or agent's file reference PF-0693 PCT	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US00/12811	International filing date (day/month/year) 10 MAY 2000	Priority date (day/month/year) 11 MAY 1999
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant INCYTE GENOMICS, INC.		

1.	This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2.	This REPORT consists of a total of <u>4</u> sheets. <input checked="" type="checkbox"/> This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT). These annexes consist of a total of <u>9</u> sheets.
3.	This report contains indications relating to the following items: <ul style="list-style-type: none"> I <input checked="" type="checkbox"/> Basis of the report II <input type="checkbox"/> Priority III <input type="checkbox"/> Non-establishment of report with regard to novelty, inventive step or industrial applicability IV <input type="checkbox"/> Lack of unity of invention V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement VI <input type="checkbox"/> Certain documents cited VII <input type="checkbox"/> Certain defects in the international application VIII <input type="checkbox"/> Certain observations on the international application

Date of submission of the demand 28 NOVEMBER 2000	Date of completion of this report 01 OCTOBER 2001
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer PREMA MERTZ
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/12811

I. Basis of the report

1. With regard to the elements of the international application: *

- ☐ the international application as originally filed
- ☒ the description:
pages _____ (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____
- ☒ the claims:
pages _____ (See Attached) _____, as originally filed
pages _____, as amended (together with any statement) under Article 19
pages _____, filed with the demand
pages _____, filed with the letter of _____
- ☒ the drawings:
pages _____ (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____
- ☒ the sequence listing part of the description:
pages _____ (See Attached) _____, as originally filed
pages _____, filed with the demand
pages _____, filed with the letter of _____

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

- ☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in printed form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. ☒ The amendments have resulted in the cancellation of:

- ☒ the description, pages _____ NONE _____
- ☒ the claims, Nos. _____ NONE _____
- ☒ the drawings, sheets/fig _____ NONE _____

5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US00/12811

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability, citations and explanations supporting such statement**1. statement**

Novelty (N)	Claims <u>1-2, 4-9, 12-95</u>	YES
	Claims <u>3, 10-11</u>	NO
Inventive Step (IS)	Claims <u>1-2, 4-9, 12-95</u>	YES
	Claims <u>3, 10-11</u>	NO
Industrial Applicability (IA)	Claims <u>1-95</u>	YES
	Claims <u>NONE</u>	NO

2. citations and explanations (Rule 70.7)

Claims 3, 10-11 lack novelty under PCT Article 33(2) as being anticipated by Robert Strausberg (1998).

Strausberg discloses a human cDNA clone isolated from a human testis cDNA library. The DNA encoding a fragment of the polypeptide of the reference, would potentially be a single amino acid. The reference discloses a polynucleotide fragment encoding a biologically active polypeptide fragment of the instant invention because a fragment of the cDNA of the reference would potentially be any nucleotide described in the instant application. Therefore, the cDNA sequence disclosed in the reference meets the limitations of a polynucleotide molecule of claims 3, 10-11 of the instant application.

Claims 1-2, 4-9, 12-95 meet the criteria set out in PCT Article 33(2)-(4), because the prior art does not teach or fairly suggest a polypeptide of amino acid sequence selected from the group consisting of SEQ ID NO:1-25, a polynucleotide encoding said polypeptide, a transgenic organism comprising said polynucleotide, an antibody to the polypeptide, a method for detecting the polynucleotide using a probe, a method of treatment with the polypeptide, a method for screening an agonist or antagonist of the polypeptide and a method of treatment with the agonist or antagonist.

----- NEW CITATIONS -----

NONE

Supplemental B x

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): C07K 14/47, 16/18; C12Q 1/68; C12N 5/10, 15/12, 15/63, 15/64; G01N 33/53, 33/567; A61K 38/16, 38/17, 48/00
and US Cl.: 530/350, 387.1, 387.9; 536/23.1, 23.5; 435/69.1, 71.1, 71.2, 325, 471, 320.1, 252.3, 254.11, 6, 7.1, 7.2;
514/2, 8, 12; 800/21

I. BASIS OF REPORT:

This report has been drawn on the basis of the description,
page(s) 1-81, as originally filed.
page(s) NONE, filed with the demand.
and additional amendments:
NONE

This report has been drawn on the basis of the claims,
page(s) 82-83, as originally filed.
page(s) NONE, as amended under Article 19.
page(s) NONE, filed with the demand.
and additional amendments:
Pages 84/1-84/9 filed with the letter of 14 August 2001.

This report has been drawn on the basis of the drawings,
page(s) NONE, as originally filed.
page(s) NONE, filed with the demand.
and additional amendments:
NONE

This report has been drawn on the basis of the sequence listing part of the description:
page(s) 1-43, as originally filed.
pages(s) NONE, filed with the demand.
and additional amendments:
NONE

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference PF-0693 PCT	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/US 00/ 12811	International filing date (day/month/year) 10/05/2000	(Earliest) Priority Date (day/month/year) 11/05/1999
Applicant INCYTE GENOMICS, INC. et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 8 sheets.
☐ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.
- ☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).
- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :
- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).
3. ☒ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

- ☐ as suggested by the applicant.
- ☐ because the applicant failed to suggest a figure.
- ☐ because this figure better characterizes the invention.

☐ None of the figures.

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US 00/12811

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 16 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the composition.
2. ☒ Claims Nos.: **18, 19, 21, 22**
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-23 (partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 18, 19, 21, 22

Present claims 18, 19, 21 and 22, directed to agonists and antagonists relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is not to be found, however, for any specific example of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, no search has been carried out for claims 18, 19, 21 and 22.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-23 partially

Polypeptide comprising SEQ ID NO:1, variants and fragments thereof, antibody binding to it; polynucleotide of SEQ ID NO:26, variants thereof, cell and transgenic organism comprising the same; probes derived from the polynucleotide and use thereof in a diagnostic method; pharmaceutical composition comprising the polypeptide and its therapeutic use; use of the polypeptide in screening assays for agonists, antagonists and compounds capable of altering the expression of the polynucleotide; therapeutic use of the agonists and antagonists.

2. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:2 and 27

3. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:3 and 28

4. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:4 and 29

5. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:5 and 30

6. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:6 and 31

7. Claims: 1-23 partially .

Idem as subject 1 for SEQ ID NOs:7 and 32

8. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:8 and 33

9. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:9 and 34

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

10. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:10 and 35

11. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:11 and 36

12. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:12 and 37

13. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:13 and 38

14. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:14 and 39

15. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:15 and 40

16. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:16 and 41

17. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:17 and 42

18. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:18 and 43

19. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:19 and 44

20. Claims: 1-23 partially

Idem as subject 1 for SEQ ID NOs:20 and 45

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

21. Claims: 1-23 partially

Idem as subject 1 for SEQ ID N0s:21 and 46

22. Claims: 1-23 partially

Idem as subject 1 for SEQ ID N0s:22 and 47

23. Claims: 1-23 partially

Idem as subject 1 for SEQ ID N0s:23 and 48

24. Claims: 1-23 partially

Idem as subject 1 for SEQ ID N0s:24 and 49

25. Claims: 1-23 partially

Idem as subject 1 for SEQ ID N0s:25 and 50

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/12811

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/78 C07K14/47 C12N15/63 A01K67/027
C07K16/18 C12Q1/68 A61K38/17 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N A01K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL [Online] Accession number AI188216, 14 October 1998 (1998-10-14) ROBERT STRAUSBERG: "qd66g12.x1 Soares testis_NHT Homo sapiens cDNA clone" XP002146658 the whole document	3,5-8, 10-14
A	WO 99 00410 A (INCYTE PHARMACEUTICALS, INC.) 7 January 1999 (1999-01-07) the whole document --- -/--	1-17,20, 23

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

6 September 2000

Date of mailing of the international search report

22.12.2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

MONTERO LOPEZ B.

INTERNATIONAL SEARCH REPORT

International Application No

P S 00/12811

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL [Online] Accession number AF151838, 1 June 1999 (1999-06-01) XP002146659 the whole document & LAI C.-H. ET AL.: "Identification of novel human genes evolutionarily conserved in Caenorhabditis elegans by comparative proteomics" GENOME RESEARCH, vol. 10, no. 5, May 2000 (2000-05), pages 703-713,</p> <p>-----</p>	<p>1-17,20, 23</p>



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/78, 14/47, C12N 15/63, A01K 67/027, C07K 16/18, C12Q 1/68, A61K 38/17, G01N 33/68		A2	(11) International Publication Number: WO 00/68380
			(43) International Publication Date: 16 November 2000 (16.11.00)
(21) International Application Number: PCT/US00/12811 (22) International Filing Date: 10 May 2000 (10.05.00) (30) Priority Data: 60/133,643 11 May 1999 (11.05.99) US 60/150,409 23 August 1999 (23.08.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/133,643 (CIP) Filed on 11 May 1999 (11.05.99) US 60/150,409 (CIP) Filed on 23 August 1999 (23.08.99) (71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive, #12, Mountain View, CA 94040 (US). TANG, Y., Tom		[CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94086 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). (74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS			
(57) Abstract			
<p>The invention provides human extracellular matrix and adhesion-associated proteins (EXMAD) and polynucleotides which identify and encode EXMAD. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of EXMAD.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of extracellular matrix and
5 adhesion-associated proteins and to the use of these sequences in the diagnosis, treatment, and
prevention of cell proliferative, immune, reproductive, neuronal, and genetic disorders.

BACKGROUND OF THE INVENTION

Extracellular Matrix Proteins

10 The extracellular matrix (ECM) is a complex network of glycoproteins, polysaccharides,
proteoglycans, and other macromolecules that are secreted from the cell into the extracellular space.
The ECM remains in close association with the cell surface and provides a supportive meshwork that
profoundly influences cell shape, motility, strength, flexibility, and adhesion. In fact, adhesion of a cell
to its surrounding matrix is required for cell survival except in the case of metastatic tumor cells, which
15 have overcome the need for cell-ECM anchorage. This phenomenon suggests that the ECM plays a
critical role in the molecular mechanisms of growth control and metastasis. (Reviewed in Ruoslahti, E.
(1996) Sci. Am. 275:72-77.) Furthermore, the ECM determines the structure and physical properties
of connective tissue and is particularly important for morphogenesis and other processes associated
with embryonic development and pattern formation.

Collagens

20 The collagens comprise a family of ECM proteins that provide structure to bone, teeth, skin,
ligaments, tendons, cartilage, blood vessels, and basement membranes. Multiple collagen proteins have
been identified. Three collagen molecules fold together in a triple helix stabilized by interchain disulfide
25 bonds. Bundles of these triple helices then associate to form fibrils. Collagen primary structure
consists of hundreds of (Gly-X-Y) repeats where about a third of the X and Y residues are Pro.
Glycines are crucial to helix formation as the bulkier amino acid side chains cannot fold into the triple
helical conformation. Because of these strict sequence requirements, mutations in collagen genes have
severe consequences. Osteogenesis imperfecta patients have brittle bones that fracture easily; in severe
30 cases patients die in utero or at birth. Ehler-Danlos syndrome patients have hyperelastic skin,
hypermobile joints, and susceptibility to aortic and intestinal rupture. Chondrodysplasia patients have
short stature and ocular disorders. Alport syndrome patients have hematuria, sensorineural deafness,
and eye lens deformation. (See Isselbacher, K.J., et al. (1994) Harrison's Principles of Internal
Medicine, McGraw-Hill, Inc., New York, NY, pp. 2105-2117; and Creighton, T.E. (1984) Proteins,

Structures and Molecular Principles, W.H. Freeman and Company, New York, NY, pp. 191-197.)

Collectins are extracellular proteins with collagen tails and globular lectin domains that play an important role in the first line immune response to microorganisms. The peripheral lectin domain permits binding to sugar residues on microorganisms, while the collagen tail interacts with phagocyte receptors or the complement system. Examples of collectins are the pulmonary surfactant proteins SP-A and SP-D (Kuroki, S.D. et al. (1998) J. Biol. Chem. 273:4783-4789).

Elastin

Elastin and related proteins confer elasticity to tissues such as skin, blood vessels, and lungs. Elastin is a highly hydrophobic protein of about 750 amino acids that is rich in proline and glycine residues. Elastin molecules are highly cross-linked, forming an extensive extracellular network of fibers and sheets. Elastin fibers are surrounded by a sheath of microfibrils which are composed of a number of glycoproteins, including fibrillin. Mutations in the gene encoding fibrillin are responsible for Marfan's syndrome, a genetic disorder characterized by defects in connective tissue. In severe cases, the aortas of afflicted individuals are prone to rupture. (Reviewed in Alberts, B., et al. (1994) Molecular Biology of the Cell, Garland Publishing, New York, NY, pp. 984-986.)

Fibronectin

Fibronectin is a large ECM glycoprotein found in all vertebrates. Fibronectin exists as a dimer of two subunits, each containing about 2,500 amino acids. Each subunit folds into a rod-like structure containing multiple domains. The domains each contain multiple repeated modules, the most common of which is the type III fibronectin repeat. The type III fibronectin repeat is about 90 amino acids in length and is also found in other ECM proteins and in some plasma membrane and cytoplasmic proteins. Furthermore, some type III fibronectin repeats contain a characteristic tripeptide consisting of Arginine-Glycine-Aspartic acid (RGD). The RGD sequence is recognized by the integrin family of cell surface receptors and is also found in other ECM proteins. Disruption of both copies of the gene encoding fibronectin causes early embryonic lethality in mice. The mutant embryos display extensive morphological defects, including defects in the formation of the notochord, somites, heart, blood vessels, neural tube, and extraembryonic structures. (Reviewed in Alberts, supra, pp. 986-987.)

Laminin

Laminin is a major glycoprotein component of the basal lamina which underlies and supports epithelial cell sheets. Laminin is one of the first ECM proteins synthesized in the developing embryo. Laminin is an 850 kilodalton protein composed of three polypeptide chains joined in the shape of a

cross by disulfide bonds. Laminin is especially important for angiogenesis and, in particular, for guiding the formation of capillaries. (Reviewed in Alberts, supra, pp. 990-991.)

Proteoglycans

- 5 There are many other types of proteinaceous ECM components, most of which can be classified as proteoglycans. Proteoglycans are composed of unbranched polysaccharide chains (glycosaminoglycans) attached to protein cores. Common proteoglycans include aggrecan, betaglycan, decorin, perlecan, serglycin, and syndecan-1. Some of these molecules not only provide mechanical support, but also bind to extracellular signaling molecules, such as fibroblast growth factor and
- 10 transforming growth factor β , suggesting a role for proteoglycans in cell-cell communication. (Reviewed in Alberts, supra, pp. 973-978.) Likewise, the glycoproteins tenascin-C and tenascin-R are expressed in developing and lesioned neural tissue and provide stimulatory and anti-adhesive (inhibitory) properties, respectively, for axonal growth (Faissner, A. (1997) Cell Tissue Res. 290:331-341).
- 15 Dentin phosphoryn (DPP) is a major component of the dentin ECM. DPP is a proteoglycan that is synthesized and expressed by odontoblasts (Gu, K., et al. (1998) Eur. J. Oral Sci. 106:1043-1047). DPP is believed to nucleate or modulate the formation of hydroxyapatite crystals. The gene encoding DPP has been mapped to human chromosome 4. Chromosome 4 contains the gene loci for two dentin genetic diseases, dentinogenesis imperfecta type II and dentin dysplasia type II (Feng, J.Q.,
- 20 et al. (1998) J. Biol. Chem. 273:9457-9464).

Mucins

- Mucins are highly glycosylated glycoproteins that are the major structural component of the mucus gel. The physiological functions of mucins are cytoprotection, mechanical protection,
- 25 maintenance of viscosity in secretions, and cellular recognition. MUC6 is a human gastric mucin that is also found in gall bladder, pancreas, seminal vesicles, and female reproductive tract (Toribara, N.W., et al. (1997) J. Biol. Chem. 272:16398-16403). The MUC6 gene has been mapped to human chromosome 11 (Toribara, N.W., et al. (1993) J. Biol. Chem. 268:5879-5885). Hemomucin is a novel
- Drosophila surface mucin that may be involved in the induction of antibacterial effector molecules
- 30 (Theopold, U., et al. (1996) J. Biol. Chem. 271:12708-12715).

Link Protein

Link protein binds to both cartilage proteoglycan and hyaluronan in cartilage ECM. This binding stabilizes the aggregation of these cartilage ECM proteins and produces supramolecular

assemblies. Link protein has been detected in other connective tissues, where it may bind proteoglycans and hyaluronan. Link protein contains a signal peptide, an immunoglobulin repeat, and link repeats (Ayad, S., et al. (1994) The Extracellular Matrix Facts Book, Academic Press, Inc., San Diego, CA, pp. 120-121).

5

Adhesion-Associated Proteins

The surface of a cell is rich in transmembrane proteoglycans, glycoproteins, glycolipids, and receptors. These macromolecules mediate adhesion with other cells and with components of the ECM. The interaction of the cell with its surroundings profoundly influences cell shape, strength, flexibility, motility, and adhesion. These dynamic properties are intimately associated with signal transduction pathways controlling cell proliferation and differentiation, tissue construction, and embryonic development.

10

Cadherins

Cadherins comprise a family of calcium-dependent glycoproteins that function in mediating cell-cell adhesion in virtually all solid tissues of multicellular organisms. These proteins share multiple repeats of a cadherin-specific motif, and the repeats form the folding units of the cadherin ECM. Cadherin molecules cooperate to form focal contacts, or adhesion plaques, between adjacent epithelial cells. The cadherin family includes the classical cadherins and protocadherins. Classical cadherins include the E-cadherin, N-cadherin, and P-cadherin subfamilies. E-cadherin is present on many types of epithelial cells and is especially important for embryonic development. P-cadherin is present on cells of the placenta and epidermis. Recent studies report that protocadherins are involved in a variety of cell-cell interactions (Suzuki, S. T. (1996) J. Cell Sci. 109:2609-2611). The intracellular anchorage of cadherins is regulated by their dynamic association with catenins, a family of cytoplasmic signal transduction proteins associated with the actin cytoskeleton. The anchorage of cadherins to the actin cytoskeleton appears to be regulated by protein tyrosine phosphorylation, and the cadherins are the target of phosphorylation-induced junctional disassembly (Aberle, H., et al. (1996) J. Cell. Biochem. 61:514-523).

15

20

25

30 Integrins

Integrins are ubiquitous transmembrane adhesion molecules that link the ECM to the internal cytoskeleton. Integrins are composed of two noncovalently associated transmembrane glycoprotein subunits called α and β . Integrins function as receptors that play a role in signal transduction. For example, binding of integrin to its extracellular ligand may stimulate changes in intracellular calcium

levels or protein kinase activity (Sjaastad, M.D. and Nelson, W.J. (1997) *BioEssays* 19:47-55). At least ten cell surface receptors of the integrin family recognize the ECM component fibronectin, which is involved in many different biological processes including cell migration and embryogenesis (Johansson, S., et al. (1997) *Front. Biosci.* 2:D126-D146).

5

Lectins

Lectins comprise a ubiquitous family of extracellular glycoproteins which bind cell surface carbohydrates specifically and reversibly, resulting in the agglutination of cells. (Reviewed in Drickamer, K. and Taylor, M.E. (1993) *Annu. Rev. Cell Biol.* 9:237-264.) This function is particularly important for activation of the immune response. Lectins mediate the agglutination and mitogenic stimulation of lymphocytes at sites of inflammation (Lasky, L.A. (1991) *J. Cell. Biochem.* 45:139-146; Palletta, E., et al. (1989) *J. Immunol.* 143:2850-2857).

Lectins are further classified into subfamilies based on carbohydrate-binding specificity and other criteria. The galectin subfamily, in particular, includes lectins that bind β -galactoside carbohydrate moieties in a thiol-dependent manner. (Reviewed in Hadari, Y.R., et al. (1998) *J. Biol. Chem.* 270:3447-3453.) Galectins are widely expressed and developmentally regulated. Because all galectins lack an N-terminal signal peptide, it is suggested that galectins are externalized through an atypical secretory mechanism. Two classes of galectins have been defined based on molecular weight and oligomerization properties. Small galectins form homodimers and are about 14-16 kilodaltons in mass, while large galectins are monomeric and about 29-37 kilodaltons.

Galectins contain a characteristic carbohydrate recognition domain (CRD). The CRD is about 140 amino acids and contains several stretches of about 1-10 amino acids which are highly conserved among all galectins. A particular 6-amino acid motif within the CRD contains conserved tryptophan and arginine residues which are critical for carbohydrate binding. The CRD of some galectins also contains cysteine residues which may be important for disulfide bond formation. Secondary structure predictions indicate that the CRD forms several β -sheets.

Galectins play a number of roles in diseases and conditions associated with cell-cell and cell-matrix interactions. For example, certain galectins associate with sites of inflammation and bind to cell surface immunoglobulin E molecules. In addition, galectins may play an important role in cancer metastasis. Galectin overexpression is correlated with the metastatic potential of cancers in humans and mice. Moreover, anti-galectin antibodies inhibit processes associated with cell transformation, such as cell aggregation and anchorage-independent growth. (See, for example, Su, Z.-Z., et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:7252-7257.)

Selectins

Selectins, or LEC-CAMs, comprise a specialized lectin subfamily involved primarily in inflammation and leukocyte adhesion. (Reviewed in Lasky, supra.) Selectins, which mediate the recruitment of leukocytes from the circulation to sites of acute inflammation, are expressed on the surface of vascular endothelial cells in response to cytokine signaling. Selectins bind to specific ligands on the leukocyte cell membrane and enable the leukocyte to adhere to and migrate along the endothelial surface. Binding of selectin to its ligand leads to polarized rearrangement of the actin cytoskeleton and stimulates signal transduction within the leukocyte (Brenner, B., et al. (1997) *Biochem. Biophys. Res. Commun.* 231:802-807; Hidari, K.I., et al. (1997) *J. Biol. Chem.* 272:28750-28756). Members of the selectin family possess three characteristic motifs: a lectin or carbohydrate recognition domain; an epidermal growth factor (EGF)-like domain; and a variable number of short consensus repeats (scr or "sushi" repeats) which are also present in complement regulatory proteins. The selectins include lymphocyte adhesion molecule-1 (LAM-1 or L-selectin), endothelial leukocyte adhesion molecule-1 (ELAM-1 or E-selectin), and granule membrane protein-140 (GMP-140 or P-selectin) (Johnston, G.I., et al. (1989) *Cell* 56:1033-1044).

Attractin

Attractin is a 134 kilodalton glycoprotein found in the serum. It is a member of the CUB family of cell adhesion proteins and binds directly to leukocytes. Attractin has a CUB domain, an EGF domain, and C-type lectin protein domains. This serum protein mediates the interaction between T lymphocytes and monocytes and leads to the adherence and spreading of monocytes that become the foci for T cell clustering. (See, Duke-Cohan, J.S., et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:11336-11341.)

Proteins Containing Leucine Rich Repeats (LRRs)

LRRs are sequence motifs, approximately 22-28 amino acids in length, found in proteins with a large variety of functions and cellular locations. Proteins containing LRRs are all thought to be involved in protein-protein interactions. The crystal structure of LRRs has been studied and found to correspond to beta-alpha structural units. These structural units form a parallel beta sheet with one surface exposed to solvent. In this way an LRR-containing protein acquires a nonglobular shape (Kobe, B. and Deisenhofer, J. (1994) *Trends Biochem. Sci.* 19:415-421). There is evidence to suggest LRRs function in signal transduction and cellular adhesion as well as in protein-protein interactions (Gay, N.J., et al. (1991) *FEBS Lett.* 29:87-91). For example, LLR proteins such as connectin and chaoptin are important cell adhesion molecules in neuronal development in Drosophila melanogaster,

and mammalian homologs are found in mouse (Taguchi, et al. (1996) Brain Res.Mol. Brain Res. 1-2:31-40).

Proteins Containing Armadillo/ β -Catenin-like Repeats

5 Various proteins such as those encoded by the Drosophila armadillo gene and the human APC gene contain amino acid repeats that interact with β -catenins. The armadillo gene is required for pattern formation within the embryonic segments and imaginal discs and is highly conserved. It is 63% identical to a human protein, plakoglobin, which is involved in adhesive junctions joining epithelial and other cells (Peifer, M. and Wieschaus, E. (1990) Cell 63:1167-1176). APC gene mutations appear to
10 initiate inherited forms of human colorectal cancer and sporadic forms of colorectal and gastric cancer (Rubinfeld, B., et al. (1993) Science 262:1731-1734). The fact that the protein encoded by APC interacts with catenin suggests a link between tumor initiation and cell adhesion (Su, L.K., et al. (1993) Science 262:1734-1737).

15 Proteins Containing C-type Lectin Domains

C-type lectin domains are found in a variety of proteins, including selectins and lecticans. Lecticans are a family of chondroitin sulfate proteoglycans that include aggrecan, versican, neurocan, and brevican. All C-type lectin proteins are involved in protein-protein interactions (Aspberg, A., et al. (1997) Proc. Natl. Acad. Sci. USA 94:10116-10121). A novel macrophage-restricted C-type lectin
20 protein has been cloned from mouse tissue. It is a type II transmembrane protein with one extracellular C-type lectin domain (Balch, S.G., et al. (1998) J. Biol. Chem. 273:18656-18664).

Bystin

Bystin is a cytoplasmic protein that binds directly to trophinin, a cell adhesion molecule, and
25 tastin. The three molecules form a complex that is involved in cell adhesion. Bystin, tastin, and trophinin are strongly expressed in cells involved in the implantation of embryos, specifically in cells at human implantation sites and in intermediate trophoblasts at the invasion front of the placenta in early pregnancy. Bystin also binds to cytokeratins. During early embryogenesis cytokeratins 8 and 18 are expressed in the trophectoderm of blastocytes. It is possible that the molecular complex formed by
30 bystin, tastin, and trophinin interacts with the cytokeratins of trophectoderm cells at the time of implantation. A key component of embryo implantation is the unique cell adhesion to endometrial epithelium that occurs and the subsequent invasion of the maternal tissue by the trophoblast. Bystin may have an important role in the signal transduction that links cell adhesion to proliferation (Suzuki, N., et al. (1998) Proc. Natl. Acad. Sci. 95:5027-5032).

Src-homology 3 (SH3) Domain-Containing Proteins

SH3 is a 60-70 amino acid motif found in a variety of signal transduction and cytoskeletal proteins. The SH3 domain is involved in mediating protein-protein interactions. Evidence suggests that the SH3 domains recognize a family of related domains or proteins in a variety of different tissues and species. One novel SH3 domain-containing protein is the 52 kilodalton focal adhesion protein (FAP52 or p52). FAP52 is localized to focal adhesions, specialized membrane domains in cultured cells that mediate the attachment of cells to the growth substratum and ECM. Focal adhesions consist of structural proteins, integrins, regulatory molecules, and signaling molecules and are involved in cell signaling. FAP52 may form part of this multimolecular complex that comprises focal adhesion sites (Merilainen, J., et al. (1997) J. Biol. Chem. 272:23278-23284).

The ECM plays an important role in cell invasive processes such as angiogenesis and tumor metastasis (Ruoslahti, supra). In particular, the glycoproteins laminin and fibronectin are implicated in the migration of tumor cells through the ECM (chemotaxis) in the course of metastasis of tumors to other tissues. The same process, chemotaxis, also promotes the migration of vascular endothelial cells to form new microvascular networks to support these tumors (tumor angiogenesis).

The discovery of new extracellular matrix and adhesion-associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative, immune, reproductive, neuronal, and genetic disorders.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, extracellular matrix and adhesion-associated proteins, referred to collectively as "EXMAD" and individually as "EXMAD-1," "EXMAD-2," "EXMAD-3," "EXMAD-4," "EXMAD-5," "EXMAD-6," "EXMAD-7," "EXMAD-8," "EXMAD-9," "EXMAD-10," "EXMAD-11," "EXMAD-12," "EXMAD-13," "EXMAD-14," "EXMAD-15," "EXMAD-16," "EXMAD-17," "EXMAD-18," "EXMAD-19," "EXMAD-20," "EXMAD-21," "EXMAD-22," "EXMAD-23," "EXMAD-24," and "EXMAD-25." In one aspect, the invention provides an isolated polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-25.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. In one alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:26-50.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, b) a naturally occurring

polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

5 Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50, c) a polynucleotide sequence complementary to a), or
10 d) a polynucleotide sequence complementary to b). The method comprises a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and b) detecting the presence or absence of said hybridization complex, and
15 optionally, if present, the amount thereof. In one alternative, the probe comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 60 contiguous nucleotides.

 The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an
20 amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional EXMAD, comprising
25 administering to a patient in need of such treatment the pharmaceutical composition.

 The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically
30 active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a

pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional EXMAD, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:26-50, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding EXMAD.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of EXMAD.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding EXMAD were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze EXMAD, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"EXMAD" refers to the amino acid sequences of substantially purified EXMAD obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of EXMAD. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of EXMAD either by directly interacting with EXMAD or by acting on components of the biological pathway in which EXMAD participates.

An "allelic variant" is an alternative form of the gene encoding EXMAD. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding EXMAD include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as EXMAD or a polypeptide with at least one functional characteristic of EXMAD. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding EXMAD, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding EXMAD. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent EXMAD. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of EXMAD is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of EXMAD. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of EXMAD either by directly interacting with EXMAD or by acting on components of the biological pathway in which EXMAD participates.

The term “antibody” refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind EXMAD polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.

Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic EXMAD, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or

amino acid sequence. The composition may comprise a dry formulation or an aqueous solution.

Compositions comprising polynucleotide sequences encoding EXMAD or fragments of EXMAD may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be
 5 deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or
 10 more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is
 15 conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
20	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
25	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
30	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
35	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide
 40 backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the

side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a
5 polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for
example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative
polynucleotide encodes a polypeptide which retains at least one biological or immunological function of
the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any
similar process that retains at least one biological or immunological function of the polypeptide from
10 which it was derived.

A "fragment" is a unique portion of EXMAD or the polynucleotide encoding EXMAD which
is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise
up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For
example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A
15 fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least
5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or
amino acid residues in length. Fragments may be preferentially selected from certain regions of a
molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino
acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown
20 in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported
by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the
present embodiments.

A fragment of SEQ ID NO:26-50 comprises a region of unique polynucleotide sequence that
specifically identifies SEQ ID NO:26-50, for example, as distinct from any other sequence in the
25 same genome. A fragment of SEQ ID NO:26-50 is useful, for example, in hybridization and
amplification technologies and in analogous methods that distinguish SEQ ID NO:26-50 from related
polynucleotide sequences. The precise length of a fragment of SEQ ID NO:26-50 and the region of
SEQ ID NO:26-50 to which the fragment corresponds are routinely determinable by one of ordinary
skill in the art based on the intended purpose for the fragment.

30 A fragment of SEQ ID NO:1-25 is encoded by a fragment of SEQ ID NO:26-50. A fragment
of SEQ ID NO:1-25 comprises a region of unique amino acid sequence that specifically identifies
SEQ ID NO:1-25. For example, a fragment of SEQ ID NO:1-25 is useful as an immunogenic peptide
for the development of antibodies that specifically recognize SEQ ID NO:1-25. The precise length of
a fragment of SEQ ID NO:1-25 and the region of SEQ ID NO:1-25 to which the fragment
35 corresponds are routinely determinable by one of ordinary skill in the art based on the intended

purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis

programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The

5 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

10 *Reward for match: 1*

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

15 *Word Size: 11*

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at

20 least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode

25 similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a

30 standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters

of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and “diagonals saved”=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the “percent similarity” between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the “BLAST 2 Sequences” tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term “humanized antibody” refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding

between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive
5 annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic
10 strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance,
20 denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a
25 similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g.,
30 paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

5 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of EXMAD which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of EXMAD which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

10 The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of EXMAD. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of EXMAD.

15 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a
20 functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which
25 comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding EXMAD, their complements, or fragments
30 thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA
35 polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid

sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary

polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.

5 This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence.
10 Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be use to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear
15 sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding EXMAD, or fragments thereof, or EXMAD itself, may comprise a bodily fluid; an
20 extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure
25 of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are
30 removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

“Substrate” refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

5 “Transformation” describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, 10 heat shock, lipofection, and particle bombardment. The term “transformed” cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A “transgenic organism,” as used herein, is any organism, including but not limited to 15 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in 20 vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, and plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention 25 into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A “variant” of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the “BLAST 2 Sequences” tool Version 2.0.9 (May-07-1999) 30 set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an “allelic” (as defined above), “splice,” “species,” or “polymorphic” variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides 35 due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may

possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human extracellular matrix and adhesion-associated proteins (EXMAD), the polynucleotides encoding EXMAD, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, immune, reproductive, neuronal, and genetic disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding EXMAD. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each EXMAD were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each EXMAD and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions

associated with nucleotide sequences encoding EXMAD. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:26-50 and to distinguish between SEQ ID NO:26-50 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express EXMAD as a fraction of total tissues expressing EXMAD. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing EXMAD as a fraction of total tissues expressing EXMAD. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding EXMAD were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:42 maps to chromosome 8 within the interval from 64.60 to 90.20 centiMorgans.

SEQ ID NO:48 maps to chromosome 2 within the interval from 193.60 to 197.60 centiMorgans.

The invention also encompasses EXMAD variants. A preferred EXMAD variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the EXMAD amino acid sequence, and which contains at least one functional or structural characteristic of EXMAD.

The invention also encompasses polynucleotides which encode EXMAD. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:26-50, which encodes EXMAD. The polynucleotide sequences of SEQ ID NO:26-50, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding EXMAD. In particular, such a variant polynucleotide sequence will have at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding EXMAD. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:26-50 which has at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:26-50. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of EXMAD.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding EXMAD, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring EXMAD, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode EXMAD and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring EXMAD under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding EXMAD or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding EXMAD and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode EXMAD and EXMAD derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding EXMAD or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:26-50 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with

machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal
cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer).
Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-
Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or
5 other systems known in the art. The resulting sequences are analyzed using a variety of algorithms
which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular
Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and
Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding EXMAD may be extended utilizing a partial nucleotide
10 sequence and employing various PCR-based methods known in the art to detect upstream sequences,
such as promoters and regulatory elements. For example, one method which may be employed,
restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic
DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.)
Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown
15 sequence from a circularized template. The template is derived from restriction fragments comprising a
known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids
Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent
to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.
(1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and
20 ligations may be used to insert an engineered double-stranded sequence into a region of unknown
sequence before performing PCR. Other methods which may be used to retrieve unknown sequences
are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).
Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo
Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in
25 finding intron/exon junctions. For all PCR-based methods, primers may be designed using
commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences,
Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a
GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to
72°C.

30 When screening for full-length cDNAs, it is preferable to use libraries that have been
size-selected to include larger cDNAs. In addition, random-primed libraries, which often include
sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library
does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5'
non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode EXMAD may be cloned in recombinant DNA molecules that direct expression of EXMAD, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express EXMAD.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter EXMAD-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of EXMAD, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby

maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding EXMAD may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser. 7*:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser. 7*:225-232.) Alternatively, EXMAD itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of EXMAD, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins. Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active EXMAD, the nucleotide sequences encoding EXMAD or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding EXMAD. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding EXMAD. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding EXMAD and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding EXMAD and appropriate transcriptional and translational

control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding EXMAD. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding EXMAD. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding EXMAD can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding EXMAD into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of EXMAD are needed, e.g. for the production of antibodies, vectors which direct high level expression of EXMAD may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of EXMAD. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of EXMAD. Transcription of sequences encoding EXMAD may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J.

6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated
5 transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding EXMAD may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader
10 sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses EXMAD in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

15 Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of
20 EXMAD in cell lines is preferred. For example, sequences encoding EXMAD can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a
25 selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase
30 genes, for use in *tk⁻* and *apr⁻* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980)

Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding EXMAD is inserted within a marker gene sequence, transformed cells containing sequences encoding EXMAD can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding EXMAD under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding EXMAD and that express EXMAD may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of EXMAD using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on EXMAD is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods. a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding EXMAD include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding EXMAD, or any fragments thereof, may be cloned into a vector

for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding EXMAD may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode EXMAD may be designed to contain signal sequences which direct secretion of EXMAD through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding EXMAD may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric EXMAD protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of EXMAD activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize

these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the EXMAD encoding sequence and the heterologous protein sequence, so that EXMAD may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled EXMAD may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of EXMAD may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of EXMAD may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of EXMAD and extracellular matrix and adhesion-associated proteins. In addition, the expression of EXMAD is closely associated with cancerous, proliferating, inflamed, nervous, reproductive, urologic, hematopoietic/immune, cardiovascular, musculoskeletal, developmental, and gastrointestinal tissues, and with cell proliferative disorders, including cancer, inflammation and the immune response. Therefore, EXMAD appears to play a role in cell proliferative, immune, reproductive, neuronal, and genetic disorders. In the treatment of disorders associated with increased EXMAD expression or activity, it is desirable to decrease the expression or activity of EXMAD. In the treatment of disorders associated with decreased EXMAD expression or activity, it is desirable to increase the expression or activity of EXMAD.

Therefore, in one embodiment, EXMAD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate,

salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an immune disorder, such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a reproductive disorder, such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; a neuronal disorder, such as akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; and a genetic disorder, such as adrenoleukodystrophy, Alport's syndrome, choroideremia, Duchenne and Becker muscular dystrophy, Down's syndrome, cystic fibrosis, chronic granulomatous disease, dentinogenesis imperfecta type II, dentin dysplasia type II, Gaucher's disease, Huntington's chorea, Marfan's syndrome, muscular dystrophy, myotonic dystrophy, pycnodysostosis, Refsum's syndrome, retinoblastoma, sickle cell anemia, thalassemia, Werner syndrome, von Willebrand's disease, Wilms' tumor, Zellweger syndrome, peroxisomal acyl-CoA oxidase deficiency, peroxisomal thiolase deficiency, peroxisomal bifunctional protein deficiency, mitochondrial carnitine palmitoyl transferase

and carnitine deficiency, mitochondrial very-long-chain acyl-CoA dehydrogenase deficiency, mitochondrial medium-chain acyl-CoA dehydrogenase deficiency, mitochondrial short-chain acyl-CoA dehydrogenase deficiency, mitochondrial electron transport flavoprotein and electron transport flavoprotein:ubiquinone oxidoreductase deficiency, mitochondrial trifunctional protein deficiency, and mitochondrial short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency.

In another embodiment, a vector capable of expressing EXMAD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified EXMAD in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of EXMAD may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of EXMAD including, but not limited to, those listed above.

In a further embodiment, an antagonist of EXMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of EXMAD. Examples of such disorders include, but are not limited to, those cell proliferative, immune, reproductive, neuronal, and genetic disorders described above. In one aspect, an antibody which specifically binds EXMAD may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express EXMAD.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding EXMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of EXMAD including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of EXMAD may be produced using methods which are generally known in the art. In particular, purified EXMAD may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind EXMAD. Antibodies to EXMAD may also be generated using methods that are well known in the art. Such antibodies may include, but are

not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with EXMAD or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to EXMAD have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of EXMAD amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to EXMAD may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce EXMAD-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in

the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for EXMAD may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between EXMAD and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering EXMAD epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for EXMAD. Affinity is expressed as an association constant, K_a, which is defined as the molar concentration of EXMAD-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple EXMAD epitopes, represents the average affinity, or avidity, of the antibodies for EXMAD. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular EXMAD epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10⁹ to 10¹² L/mole are preferred for use in immunoassays in which the EXMAD-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10⁶ to 10⁷ L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of EXMAD, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of EXMAD-antibody

complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

5 In another embodiment of the invention, the polynucleotides encoding EXMAD, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding EXMAD may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding EXMAD. Thus, complementary molecules or fragments may be used to modulate EXMAD activity, or to achieve regulation of gene function. Such technology
10 is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding EXMAD.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to
15 construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding EXMAD. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding EXMAD can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding EXMAD. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the
20 absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing
25 complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding EXMAD. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases,
30 transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding EXMAD.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding EXMAD. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such

therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of EXMAD, antibodies to EXMAD, and mimetics, agonists, antagonists, or inhibitors of EXMAD. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene

glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of EXMAD, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the

active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs.

- 5 An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

- 10 A therapeutically effective dose refers to that amount of active ingredient, for example EXMAD or fragments thereof, antibodies of EXMAD, and agonists, antagonists or inhibitors of EXMAD, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

- 20 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy.
- 25 Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

- 30 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind EXMAD may be used for the

diagnosis of disorders characterized by expression of EXMAD, or in assays to monitor patients being treated with EXMAD or agonists, antagonists, or inhibitors of EXMAD. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics.

Diagnostic assays for EXMAD include methods which utilize the antibody and a label to detect

- 5 EXMAD in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

- A variety of protocols for measuring EXMAD, including ELISAs, RIAs, and FACS, are
10 known in the art and provide a basis for diagnosing altered or abnormal levels of EXMAD expression. Normal or standard values for EXMAD expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to EXMAD under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of EXMAD expressed
15 in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

- In another embodiment of the invention, the polynucleotides encoding EXMAD may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and
20 quantify gene expression in biopsied tissues in which expression of EXMAD may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of EXMAD, and to monitor regulation of EXMAD levels during therapeutic intervention.

- In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding EXMAD or closely related molecules may be used
25 to identify nucleic acid sequences which encode EXMAD. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding EXMAD, allelic variants, or related sequences.

- 30 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the EXMAD encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:26-50 or from genomic sequences including promoters, enhancers, and introns of the EXMAD gene.

Means for producing specific hybridization probes for DNAs encoding EXMAD include the

cloning of polynucleotide sequences encoding EXMAD or EXMAD derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding EXMAD may be used for the diagnosis of disorders associated with expression of EXMAD. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an immune disorder, such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a reproductive disorder, such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and

galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; a neuronal disorder, such as akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; and a genetic disorder, such as adrenoleukodystrophy, Alport's syndrome, choroideremia, Duchenne and Becker muscular dystrophy, Down's syndrome, cystic fibrosis, chronic granulomatous disease, dentinogenesis imperfecta type II, dentin dysplasia type II, Gaucher's disease, Huntington's chorea, Marfan's syndrome, muscular dystrophy, myotonic dystrophy, pycnodysostosis, Refsum's syndrome, retinoblastoma, sickle cell anemia, thalassemia, Werner syndrome, von Willebrand's disease, Wilms' tumor, Zellweger syndrome, peroxisomal acyl-CoA oxidase deficiency, peroxisomal thiolase deficiency, peroxisomal bifunctional protein deficiency, mitochondrial carnitine palmitoyl transferase and carnitine deficiency, mitochondrial very-long-chain acyl-CoA dehydrogenase deficiency, mitochondrial medium-chain acyl-CoA dehydrogenase deficiency, mitochondrial short-chain acyl-CoA dehydrogenase deficiency, mitochondrial electron transport flavoprotein and electron transport flavoprotein:ubiquinone oxidoreductase deficiency, mitochondrial trifunctional protein deficiency, and mitochondrial short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency. The polynucleotide sequences encoding EXMAD may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered EXMAD expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding EXMAD may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding EXMAD may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding EXMAD in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of EXMAD, a normal or standard profile for expression is established. This may be accomplished by

combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding EXMAD, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding EXMAD may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding EXMAD, or a fragment of a polynucleotide complementary to the polynucleotide encoding EXMAD, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of EXMAD include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic

variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding EXMAD may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding EXMAD on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or

affected individuals.

In another embodiment of the invention, EXMAD, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between EXMAD and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with EXMAD, or fragments thereof, and washed. Bound EXMAD is then detected by methods well known in the art. Purified EXMAD can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding EXMAD specifically compete with a test compound for binding EXMAD. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with EXMAD.

In additional embodiments, the nucleotide sequences which encode EXMAD may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No.60/133,643 and U.S. Ser. No.60/150,409 are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic

solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUEScript plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-

well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

5 cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as
10 the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the
15 cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions,
20 references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between
25 two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

30 The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programing, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation

using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:26-50. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding EXMAD occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous,

reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table

3.

V. Chromosomal Mapping of EXMAD Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:40-50 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:40-50 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

The genetic map locations of SEQ ID NO:42 and SEQ ID NO:48 are described in The Invention as ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters.

VI. Extension of EXMAD Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:26-50 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR

was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE

Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:26-50 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

5 VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:26-50 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

20 VIII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or

fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

IX. Complementary Polynucleotides

Sequences complementary to the EXMAD-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring EXMAD. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of EXMAD. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the EXMAD-encoding transcript.

X. Expression of EXMAD

Expression and purification of EXMAD is achieved using bacterial or virus-based expression systems. For expression of EXMAD in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express EXMAD upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of EXMAD in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding EXMAD by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, EXMAD is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from EXMAD at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified EXMAD obtained by these methods can be used directly in the following activity assay.

XI. Demonstration of EXMAD Activity

An assay for EXMAD activity measures the disruption of cytoskeletal filament networks upon overexpression of EXMAD in cultured cell lines. (Reznicek, G. A. et al. (1998) J. Cell Biol. 141:209-225.) cDNA encoding EXMAD is subcloned into a mammalian expression vector that drives high levels of cDNA expression. This construct is transfected into cultured cells, such as rat kangaroo PtK2 or rat bladder carcinoma 804G cells. Actin filaments and intermediate filaments such as keratin and vimentin are visualized by immunofluorescence microscopy using antibodies and techniques well known in the art. The configuration and abundance of cytoskeletal filaments can be assessed and quantified using confocal imaging techniques. In particular, the bundling and collapse of cytoskeletal filament networks are indicative of EXMAD activity.

Alternatively, an assay for EXMAD activity measures the amount of cell aggregation induced by overexpression of EXMAD. In this assay, cultured cells such as NIH3T3 are transfected with cDNA encoding EXMAD contained within a suitable mammalian expression vector under control of a strong promoter. Cotransfection with cDNA encoding a fluorescent marker protein, such as Green Fluorescent Protein (Clontech), is useful for identifying stable transfectants. The amount of cell agglutination, or clumping, associated with transfected cells is compared with that associated with untransfected cells. The amount of cell agglutination is a direct measure of EXMAD activity.

Alternatively, cell adhesion activity in EXMAD is measured in a 96-well plate assay in which wells are first coated with EXMAD by adding solutions of EXMAD of varying concentrations to the wells. Excess EXMAD is washed off with saline, and the wells incubated with a solution of 1% bovine serum albumin to block non-specific cell binding. Aliquots of a cell suspension of a suitable cell type are then added to the wells and incubated for a period of time at 37 °C. Non-adhered cells are washed

off with saline and the cells stained with a suitable cell stain such as Coomassie blue. The intensity of staining is measured using a variable wavelength 96-well plate reader and compared to a standard curve to determine the number of cells adhering to the EXMAD coated plates. The degree of cell staining is proportional to the cell adhesion activity of EXMAD in the sample.

Alternatively, EXMAD activity is also measured by the interaction of EXMAD with other molecules. EXMAD, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) *Biochem. J.* 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled EXMAD, washed, and any wells with labeled EXMAD complex are assayed. Data obtained using different concentrations of EXMAD are used to calculate values for the number, affinity, and association of EXMAD with the candidate molecules.

XII. Functional Assays

EXMAD function is assessed by expressing the sequences encoding EXMAD at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 μg of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μg of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) *Flow Cytometry*, Oxford, New York NY.

The influence of EXMAD on gene expression can be assessed using highly purified

populations of cells transfected with sequences encoding EXMAD and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding EXMAD and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIII. Production of EXMAD Specific Antibodies

EXMAD substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the EXMAD amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-EXMAD activity by, for example, binding the peptide or EXMAD to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIV. Purification of Naturally Occurring EXMAD Using Specific Antibodies

Naturally occurring or recombinant EXMAD is substantially purified by immunoaffinity chromatography using antibodies specific for EXMAD. An immunoaffinity column is constructed by covalently coupling anti-EXMAD antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing EXMAD are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of EXMAD (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/EXMAD binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and EXMAD is collected.

XV. Identification of Molecules Which Interact with EXMAD

EXMAD, or biologically active fragments thereof, are labeled with ^{125}I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled EXMAD, washed, and any wells with labeled EXMAD complex are assayed. Data obtained using different concentrations of EXMAD are used to calculate values for the number, affinity, and association of EXMAD with the candidate molecules.

Alternatively, molecules interacting with EXMAD are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, *Nature* 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments.

Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	26	398269	PITUNOT02	265928H1 (HNT2AGT01), 398269H1 and 398269R6 (PITUNOT02), 516201R6 (MMLR1DT01), 822473R6 (KERANOT02), 1265919F1 (BRAINOT09), 1356244F6 (LUNGNOT09), 1379344T6 (LUNGNOT10), 3586102H1 (293TF4T01), SBLA02091F1, SBLA01281F1
2	27	1258888	MENITUT03	125888H1 (MENITUT03), 1373184H1 (BSTMN02), 2420735R6 (SCORN02), 2697827F3 (UTRSNOT12), 2990569T6 (KIDNFET02), SBGA02402F1, SBGA05599F1, SBGA01330F1, SBGA07058F3
3	28	1375891	LUNGNOT10	1375891H1 (LUNGNOT10), 2251462R6 (OVARUT01), 4542640H1 (THYRTMT01), SAXA00188F1, SAXA00819F1, SAXA00256F1, SAXA00101F1, SZAC00197F1
4	29	1524355	UCMCL5T01	008503T6 (HMCINOT01), 425033R6 (BLADNOT01), 1299403T6 (BRSTNOT07), 1524355H1 (UCMCL5T01), 2480893F6 (SMCANOT01), 3072568F6 (UTRSNOT01), 3077770H1 (BONEUNT01), 3521659H1 (LUNGNOT03), 3810130H1 (CONTTUT01), 4187444H1 (BRSTNOT31)
5	30	1598937	BLADNOT03	307298R6 (HEARNOT01), 637901F1 (BRSTNOT03), 872833R1 (LUNGAST01), 1360462F1 (LUNGNOT12), 1598937H1 (BLADNOT03), 1688334H1 (PROSTUT10), 2048691F6 (LIVRFET02), 3604769H1 (LUNGNOT30)
6	31	1725801	PROSNOT14	359107F1 and 359107R1 (SYNORAB01), 1725801H1 and 1725801X18C1 (PROSNOT14), 2853280H1 (CONNNOT02), SBWA02129V1
7	32	1730482	BRSTTUT08	1261313R1 (SYNORAT05), 1321141F1 (BLADNOT04), 1484641F1 (CORPNOT02), 1730482H1 (BRSTTUT08), 1848053F6 (OVARNOT07), 2208990F6 (SINTFFET03), 2691973F6 (LUNGNOT23), 2811183H1 (OVARNOT10), 3097712H1 (CERVNOT03), 3110665H1 (BRSTNOT17), 3738668H1 (MENTNOT01)
8	33	1810058	PROSTUT12	571697H1 (OVARNON01), 1704596F6 (DUODNOT02), 1810058H1, 1810548F6, and 1810548T6 (PROSTUT12)
9	34	2040679	HIPONON02	2040679H1 and 2040679R6 (HIPONON02), 2380160F6 (ISLTNOT01), 2621171T6 (KERANOT02), 2869976F6 (THYRNOT10)
10	35	2960051	ADRENOT09	2960051F6 and 2960051H1 (ADRENOT09), SBVA05142V1, SBVA03774V1, SBVA03935V1

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
11	36	3117318	LUNGTUT13	393775H1 (TMLR2DT01), 486988H1 (HNT2AGT01), 3117318F6 and 3117318H1 (LUNGTUT13), 3293662F6 (TLYJINT01), SBMA01131F1
12	37	3486992	EPIGNOT01	2615184H1 (GBLANOT01), 3486992H1 (EPIGNOT01), SBKA01303F1.comp, SBKA03723F1.comp, SBKA02206F1, SBKA01625F1.comp, SBKA02769F1, SBKA03712F1, SBKA02365F1, SBKA01975F1
13	38	4568384	HELATXT01	080350F1 (SYNORAB01), 320872H1 (EOSIHET02), 1418995F1 (KIDNNOT09), 1473647T1 (LUNGTUT03), 1664971F6 (BRSTNOT09), 1738547F6 (COLNNOT22), 2367046F6 (ADRENOT07), 4568384F6 and 4568384H1 (HELATXT01)
14	39	4586187	OVARNOT13	306792F1 and 306792X11R1 (HEARNOT01), 632244F1 (KIDNNOT05), 876626R1 (LUNGAST01), 2451238F6 (ENDANOT01), 2881494F6 (UTRSTUT05), 4586187H1 (OVARNOT13), 5852878H1 (FIBAUNT02), SZZZ01051R1
15	40	401801	TMLR3DT01	401801T6 and 401801H1 (TMLR3DT01), 938106H1 (CERVNOT01), 2603123T6 (UTRSNOT10), 2607556H1 (LUNGTUT07)
16	41	1721842	BLADNOT06	1721842H1, 1721842F6 and 1721842T6 (BLADNOT06), 2010387R6 (TESTNOT03), 4884119H1 (LUNLTMT01)
17	42	1833221	BRAINON01	001593H1 (U937NOT01), 389513R1 (THYMNOT02), 428370R6 (BLADNOT01), 493657H1 (HNT2NOT01), 1263824R1 (SYNORAT05), 1833221H1 (BRAINON01), 1907733F6 (CONNTUT01), 1997529R6 (BRSTTUT03), 2174658F6 (ENDCNOT03), 3114306H1 (BRSTNOT17), 3233178H1 (COLNUCT03), 4788994F6 (EPIBUNT01), 5541215H1
18	43	2041168	HIPONON02	849897R1 (NGANNOT01), 908128R2 (COLNNOT09), 999830R6 (KIDNTUT01), 1639572T6 (UTRSNOT06), 1686825F6 (PROSNOT15), 2041168H1 (HIPONON02), 2582551H1 (KIDNTUT13), 2867048H1 (KIDNNOT20), 3226063F6 (TLYJINT01), 3226063H1 (TLYJINT01), 3466031H1 (293TF2T01), 4662252H2 (BRSTTUT20), SBIA03151D1
19	44	2365794	ADRENOT07	874804H1 (LUNGAST01), 1318960T1 (BLADNOT04)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
20	45	2618452	GBLANOT01	1730514F6 (BRSTTUT08), 2225286F6 (SEMVNOT01), 2225720F6 (SEMVNOT01), 2618452F6 and 2618452H1 (GBLANOT01), 2618457F6 (GBLANOT01), 3248134H1 (SEMVNOT03), 3250560H1 (SEMVNOT03), 3538176F6 (SEMVNOT04), 4068913H1 (SEMVNOT05)
21	46	2622288	KERANOT02	223636F1 (PANCNOT01), 490914R6 (HNT2AGT01), 530368R6 (BRAINOT03), 850583R1 (NGANNOT01), 898618R1 (BRSTTUT03), 932484R6 (CERVNOT01), 1302418F1 (PLACNOT02), 1368735R1 (SCORN02), 1486177F6 (CORPNOT02), 1726367F6 (PROSNOT14), 2516869H1 (LIVRTUT04), 2622288R6 and 2622288H2 (KERANOT02), 3043955H1 (HEAANOT01), 3398316H1 (UTRSNOT16), 3938796H1 (SKINBIT01), 4043471H1 (LUNGNOT35)
22	47	2806595	BLADTUT08	643445R6 (BRSTTUT02), 2806595F6 and 2806595H1 (BLADTUT08), SBRA04014D1, SBRA03510D1
23	48	2850987	BRSTTUT13	1300925F1 (BRSTNOT07), 1339833F1 (COLNTUT03), 1347463F6 (PROSNOT11), 1347463T6 (PROSNOT11), 1899642F6 (BLADTUT06), 2715093F6 (THYRN09), 2726463F6 (OVRTUT05), 2850987H1 (BRSTTUT13), 2893008H1 (LUNGFET04), 3336701F6 (SPLNNOT10), 3341661H1 (SPLNNOT09), SXAF00652V1, SXAF03272V1
24	49	3557211	LUNGNOT31	958552H1 (KIDNNOT05), 2953281F6 and 2953281T6 (KIDNFET01), 3557211F6 and 3557211H1 (LUNGNOT31), 4306204H1 (GBLADIT01), 4420950F6 (LIVRDIT02) g2188176, g1424165
25	50	4675668	NOSEDIT02	1519431T6 (BLADTUT04), 2447058F6 (THP1NOT03), 2758306R6 (THP1AZS08), 2758306T6 (THP1AZS08), 3589494H1 (293TF5T01), 3813434H1 (TONSNOT03), 4675668H1 (NOSEDIT02), 5175727H1 (EP1BTXT01), 5313381H1 (KIDETXS02)

Table 2

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
1	309	T153 S29 S140 T153 S162 T168 S233 S258 T285 S290 T87 T159 T265	N108 N305	Signal peptide: M1-A31	similar to B. Subtilis surfactin (SFP) protein g3880360	BLAST SPSCAN
2	554	S57 S146 S265 T275 S389 T495 T496 S497 S551 S25 S34 T87 S115 S180 S212 S242 S289 T308 S361 T388 T504	N398	EGF-like domain: C98-C132 C138-C172 C178-C217 C223-C258 Cell adhesion: R363-D365 Signal peptide: M1-G21	fibulin-2 [Mus musculus] g437047	BLAST PRINTS BLOCKS PFAM MOTIFS SPSCAN HMM
3	482	S87 T37 T108 T131 S133 S148 T165 T246 S254 T256 S269 S283 S333 S404 T414 T431 S28 T29 S65 T335 T431 S446 S460 T464	N252 N445 N451	Signal peptide: M1-G22	gastric mucin [Sus scrofa] g915208	BLAST MOTIFS SPSCAN HMM
4	735	S506 S153 S243 T259 S304 T317 T378 S414 T502 S575 S670 S688 S698 S44 T116 S258 S324 S350 S356 S396 T437 T515 S610 S620 Y53	N70 N97 N144 N188 N412	Kelch motif: T284-K330 C469-G513	muskelin [Mus musculus] g3493462	BLAST PFAM

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
5	424	T209 S256 S276 T86 S311 S319 T347 S15 S354 S394 S107 Y53 S153 T217 S258 S408 T306 S358 S383		SH3 domain: V366-V422	Focal adhesion protein (FAP52) [Gallus gallus] g2217964	BLAST PFAM PRINTS BLOCKS
6	420	S293 T63 T73 S99 S101 S222 T359 T48 T63 S138 T159 S406 S409 Y53	N79 N205	Signal peptide: M1-L29 EGF-like domain: T174-C192 Cysteine-rich pattern: C181-C192	HT protein [Cricetulus griseus] g1216486	BLAST PRINTS SPSCAN MOTIFS HMM
7	795	S41 T94 S145 S243 T297 S442 S451 T687 S103 T111 T129 S184 T428 S647	N383 N387	Cell adhesion: R606-D608 von Willebrand factor type A domain: D31-L204 transmembrane domain: I50-T77	collagen type XIV [Homo sapiens] g2065167	BLAST MOTIFS PFAM PRINTS HMM
8	306	T69 T133 S255 T279 T22		Signal peptide: M1-S19 Clq domain: G149-P175 A203-I226 H227-L302	saccular collagen [Lepomis macrochirus] g687606	BLAST PFAM PRINTS BLOCKS SPSCAN HMM

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
9	338	S5 S53 S66 T119 T246 S23 T65 S102 S151 S251 T277	N217 N332	Signal peptide: M1-S22 Leucine-rich repeats domain: S102-T147 S151-I196 N197-A243	LRR47 [Drosophila melanogaster] g415947	BLAST PFAM PRINTS SPSCAN HMM
10	164	S42 S75 T160 S44 S49		Signal peptide: M1-G20 von Willebrand factor C-type domain: C103-C157	extracellular matrix protein [Homo sapiens] g3786312	BLAST PFAM BLOCKS SPSCAN HMM
11	327	S292 S30 S35 S63 T92 T14 T102 T179 S198 T285	N54 N61 N75 N85 N100 N189 N196 N213 N218 N322	Signal peptide: M1-P29 Ig domain: P81-F144 G173-A239 Transmembrane domain: V254-A276	embigin protein [Rattus norvegicus] g3355709	BLAST PFAM SPSCAN HMM
12	716	S21 T49 T54 T87 T98 S245 T315 T471 T519 T590 S624 S692 T705 S176 S384 S473 S494 T513 S542 T560 T571 T605 T613 S664 T709 Y581	N69 N96 N106 N117 N385 N517 N582 N611	Signal peptide: M1-S25 Leucine-rich repeats domain: N96-S143 N192-D239 S240-L287 R288-P337 A338-N385 Transmembrane domain: M639-F656	leucine-rich- repeat protein [Mus musculus] g1228052	BLAST PFAM PRINTS SPSCAN HMM

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
13	665	T147 S45 S86 S110 S121 T147 S160 T200 S205 S225 S247 S299 S301 S309 S335 S336 S341 S343 T386 S388 T400 T448 S506 S534 S545 S580 S581 S582 S597 S602 S615 S23 S82 S100 S162 S183 T199 S217 S221 S329 S347 T429 T501 T558 T563 T608 Y445 Y559	N119 N242 N424 N427 N634		50kDa lectin [Bombyx mori] g500858	BLAST
14	547	T60 S31 T87 T175 S213 T357 T452 T474 S476 T488 S203 T420 Y424	N15 N76 N85 N104 N128 N154 N191 N221 N242 N418	Lectin C-type domain: L473-C535 T488-L547 Cell adhesion: R256-D258	CSR1 (cellular stress response protein) [Homo sapiens] g6230372	BLAST PFAM BLOCKS MOTIFS PPROFILES SCAN
15	109	S85 S38	N22		Attractin; DPPT- L [Homo sapiens] g3676347	BLAST-GenBank MOTIFS
16	192	S10 S87 T92 T157 T165 T170 S19 S46	N8 N103	Leucine Rich Repeat Domain: L81-I94 L126-M139		BLIMPS- PRINTS MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
17	575	T150 S171 S299 S85 S98 S117 S118 S126 S142 S170 S203 S237 S239 S333 S415 S467 T473 S524 T557 S558 S562 S32 S92 S104 S128 S134 T149 T150 S167 S188 S260 S270 S280 S289 S389 S536	N68 N96 N234 N366 N569		axotrophin [Mus musculus] g5052031 dentin phosphoryn [Homo sapiens] g4322670	BLAST-GenBank MOTIFS
18	342	S73 S24 S82 S207 S315 S96 T176	N31 N152 N180 N193	Armadillo/beta- catenin-like repeats: A104-A113		BLIMPS-PFAM MOTIFS
19	110	S80		Signal Peptide: M1-G45 Transmembrane Domain: G48-G71 G91-Y110 Legume lectins signature: V4-F54		SPSCAN HMMER PROFILES SCAN MOTIFS
20	571	S482 T502 T11 T40 S88 T180 S231 T339 T383 T402 T409 T436 T447 S482 T491	N66 N229 N434 N498	Mucin domain: P101 - S430 Cystine knot domain: C481-C569	mucin [Homo sapiens] g292046	BLAST-GenBank BLAST-DOMO HMMER-PFAM MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
21	262	S69 S146 S172 S41 T54 T59 T101 T102 T107 Y170		Signal Peptide: M1-G25	single-pass transmembrane protein [Rattus norvegicus] g6978944 antigen [Homo sapiens] g188543	SPSCAN HMMER MOTIFS BLAST-GenBank
22	172	S29 T53 S111 S80 Y144		Signal Peptide: M1-G17 Protein proteoglycan core glycoprotein precursor cartilage repeat lectin Ig fold : G63-I149 Immunoglobulin: E52-S156	link protein [Mus musculus] g4218976	BLAST-GenBank BLAST-PRODOM BLAST-DOMO SPSCAN HMMER MOTIFS
23	571	S16 T36 T294 S396 S403 T445 S23 T176 S487	N100 N174 N434 N567	Mitochondrial energy transfer proteins signature: P404-F412 Transmembrane domains: T94-K116 F520-F539 L58-I78 I341-W362 I375-M393 I453-F472 Laminin b: Y538-K558	cell adhesion regulator [Rattus norvegicus] g4098299	BLAST- GenBank, HMMER-PFAM HMMER MOTIFS

Table 2 (cont.)

Protein SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequence	Homologous Sequence	Analytical Methods
24	455	S18 S23 S143 S270 S81 T186 S196 T208 S230 T240 T256 S418 S452 Y223	N138 N217 N288	Signal peptidases I signature: G43-F50 Lectin c-type: C329-S452 Cell attachment sequence: R183-D185	lectin BRA-3 [Megabalanus rosa] g407227	BLAST-GenBank BLAST-DOMO HMMER-PFAM MOTIFS
25	437	S98 T146 T160 S211 T220 T301 S55 T86 T156 S197 T369 Y265 Y334 Y350		ENP1 protein nuclear protein: E157-D431	bystin [Mus musculus] g2738509	BLAST-GenBank BLAST-PRODOM MOTIFS

Table 3

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
26	242-286	Nervous (0.264) Reproductive (0.198)	Cancer (0.462) Cell proliferation (0.242) Inflammation (0.176)	PSPORT
27	272-316	Nervous (0.438) Reproductive (0.188) Developmental (0.188)	Cancer (0.438) Cell proliferation (0.250) Inflammation (0.188)	pINCY
28	218-262	Gastrointestinal (0.244) Nervous (0.195) Reproductive (0.171)	Cancer (0.488) Inflammation (0.195) Cell proliferation (0.146)	pINCY
29	488-532 1082-1126	Reproductive (0.265) Nervous (0.206) Hematopoietic/immune (0.147)	Cancer (0.500) Cell proliferation (0.324) Inflammation (0.235)	PBLUESCRIPT
30	542-586	Reproductive (0.321) Cardiovascular (0.143) Musculoskeletal (0.143)	Cancer (0.500) Inflammation (0.107) Cell proliferation (0.107)	pINCY
31	217-261	Nervous (0.265) Reproductive (0.253) Cardiovascular (0.108)	Cancer (0.482) Inflammation (0.145) Cell proliferation (0.145)	pINCY
32	36-80	Reproductive (0.333) Gastrointestinal (0.154) Developmental (0.115)	Cancer (0.462) Inflammation (0.167) Cell proliferation (0.154)	pINCY
33	218-262	Reproductive (0.571) Gastrointestinal (0.286) Cardiovascular (0.143)	Trauma (0.286) Cancer (0.143) Inflammation (0.143)	pINCY
34	111-155	Gastrointestinal (0.364) Nervous (0.182) Cardiovascular (0.091)	Cancer (0.364) Inflammation (0.273) Cell proliferation (0.182)	PSPORT
35	271-315	Musculoskeletal (0.286) Reproductive (0.286) Cardiovascular (0.143)	Cancer (0.286) Inflammation (0.143) Neurological (0.143)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
36	542-586 866-910	Hematopoietic/Immune (0.526) Reproductive (0.158) Nervous (0.105)	Cancer (0.368) Inflammation (0.474) Cell proliferation (0.158)	pINCY
37	811-855	Nervous (0.267) Reproductive (0.267) Musculoskeletal (0.133)	Cancer (0.600) Inflammation (0.200) Cell proliferation (0.133)	pINCY
38	380-424 974-1018	Reproductive (0.200) Gastrointestinal (0.164) Nervous (0.145)	Cancer (0.436) Cell proliferation (0.309) Inflammation (0.200)	pINCY
39	434-479 975-1019	Reproductive (0.296) Cardiovascular (0.259) Hematopoietic/Immune (0.111)	Cancer (0.315) Inflammation (0.204) Trauma (0.204)	pINCY
40	555-614	Cardiovascular (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (0.667) Cancer (0.333)	PBLUESCRIPT
41	743-802	Nervous (0.353) Reproductive (0.176) Urologic (0.176)	Cancer (0.471) Inflammation (0.411) Cell Proliferation (0.118)	pINCY
42	429-488 1029-1088	Reproductive (0.213) Nervous (0.191) Cardiovascular (0.169)	Cancer (0.472) Inflammation (0.394) Cell Proliferation (0.180)	PSPORT1
43	967-1026	Nervous (0.228) Reproductive (0.213) Gastrointestinal (0.110)	Cancer (0.504) Inflammation (0.291) Cell Proliferation (0.197)	PSPORT1
44	164-223	Reproductive (0.241) Cardiovascular (0.167) Gastrointestinal (0.148)	Cancer (0.481) Inflammation (0.315) Cell Proliferation (0.167)	pINCY
45	110-169	Gastrointestinal (0.562) Reproductive (0.312) Nervous (0.062) Urologic (0.062)	Cancer (0.500) Inflammation (0.312) Cell Proliferation (0.062)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Unique Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
46	273-332 759-818	Nervous (0.347)	Cancer (0.430)	PSPORT1
		Reproductive (0.223)	Inflammation (0.364)	
		Cardiovascular (0.132)	Cell Proliferation (0.124)	
47	218-277	Gastrointestinal (0.200)	Cancer (0.533)	pINCY
		Nervous (0.200)	Inflammation (0.334)	
		Reproductive (0.200)	Cell Proliferation (0.133)	
48	341-400	Reproductive (0.294)	Cancer (0.476)	pINCY
		Gastrointestinal (0.168)	Inflammation (0.329)	
		Cardiovascular (0.126)	Cell Proliferation (0.168)	
49	266-325 542-601	Developmental (0.277)	Cell Proliferation (0.444)	pINCY
		Gastrointestinal (0.222)	Inflammation (0.444)	
		Nervous (0.167) Urologic (0.167)	Cancer (0.167)	
50	165-224	Hematopoietic/Immune (0.216)	Cancer (0.568)	pINCY
		Reproductive (0.216)	Cell Proliferation (0.324)	
		Gastrointestinal (0.135)	Inflammation (0.297)	

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
26	PITUNOT02	The library was constructed using RNA obtained from Clontech. The RNA was isolated from the pituitary glands removed from a pool of 87 male and female donors, 15 to 75 years old.
27	MENITUT03	The library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
28	LUNGNOT10	The library was constructed using RNA isolated from the lung tissue of a Caucasian male fetus, who died at 23 weeks' gestation.
29	UCMCL5T01	The UCMCL5T01 library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of 12 individuals. The cells were cultured for 12 days with IL-5 before RNA was obtained from the pooled lysates.
30	BLADNOT03	The library was constructed using RNA isolated from the bladder tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology for the associated tumor tissue indicated grade 3 invasive transitional cell carcinoma. Patient history included malignant neoplasm of the uterus, atherosclerosis, and atrial fibrillation. Family history included acute renal failure, osteoarthritis, and atherosclerosis.
31	PROSNOT14	The library was constructed using RNA isolated from diseased prostate tissue removed from a 60-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+4). The patient presented with elevated prostate specific antigen (PSA). Patient history included a kidney cyst and hematuria. Family history included benign hypertension, cerebrovascular disease, and arteriosclerotic coronary artery disease.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
32	BRSTTUT08	The library was constructed using RNA isolated from breast tumor tissue removed from a 45-year-old Caucasian female during unilateral extended simple mastectomy. Pathology indicated invasive nuclear grade 2-3 adenocarcinoma, ductal type, with 3 of 23 lymph nodes positive for metastatic disease. Greater than 50% of the tumor volume was in situ, both comedo and non-comedo types. Immunostains were positive for estrogen/progesterone receptors, and uninvolved tissue showed proliferative changes. The patient concurrently underwent a total abdominal hysterectomy. Patient history included valvuloplasty of mitral valve without replacement, rheumatic mitral insufficiency, and rheumatic heart disease. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
33	PROSTUT12	The library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
34	HIPONON02	This normalized hippocampus library was constructed from 1.13M independent clones from a normal hippocampus library. RNA was isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis. The normalization and hybridization conditions were adapted from Soares et al. (Proc.Natl.Acad.Sci. USA (1994) 91:9228).
35	ADRENOT09	The library was constructed using RNA isolated from left adrenal gland tissue removed from a 43-year-old Caucasian male during nephroureterectomy, regional lymph node excision, and unilateral left adrenalectomy. Pathology for the associated tumor tissue indicated a grade 2 renal cell carcinoma mass in the posterior lower pole of the left kidney with invasion into the renal pelvis.
36	LUNGUT13	The library was constructed using RNA isolated from tumorous lung tissue removed from the right upper lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology indicated invasive grade 3 (of 4) adenocarcinoma. Family history included atherosclerotic coronary artery disease, and type II diabetes.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
37	EPIGN0T01	The library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
38	HELATXT01	The library was constructed using RNA isolated from HeLa cells treated with TNF- α and IL-1 β , 10ng/nl each for 20 hours. The HeLa cell line is derived from cervical adenocarcinoma removed from a 31-year-old Black female.
39	OVARNOT13	The library was constructed using RNA isolated from left ovary tissue removed from a 47-year-old Caucasian female during a vaginal hysterectomy with bilateral salpingo-oophorectomy, and dilation and curettage. Pathology for the associated tumor tissue indicated a single intramural leiomyoma. The endometrium was in the secretory phase. The patient presented with metrorrhagia. Patient history included hyperlipidemia and benign hypertension. Family history included colon cancer, benign hypertension, atherosclerotic coronary artery disease, and breast cancer.
40	TMLR3DT01	Library was constructed using RNA isolated from non-adherent and adherent peripheral blood mononuclear cells collected from two unrelated Caucasian male donors (25 and 29 years old). Cells from each donor were purified on Ficoll Hypaque, then co-cultured for 96 hours in medium containing normal human serum at a cell density of 2×10^6 cells/ml. The non-adherent and adherent cell populations were pooled, washed once in PBS, lysed in a buffer containing GuSCN, and spun through CsCl to obtain RNA.
41	BLADNOT06	Library was constructed using RNA isolated from the posterior wall bladder tissue removed from a 66-year-old Caucasian male during a radical prostatectomy, radical cystectomy and urinary diversion. Pathology for the associated tumor tissue indicated grade 3 transitional cell carcinoma on the anterior wall of the bladder and urothelium. Patient history included lung neoplasm, and tobacco abuse in remission. Family history included a malignant breast neoplasm, tuberculosis, cerebrovascular disease, atherosclerotic coronary artery disease, and lung cancer.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
42	BRAINON01	Library was constructed and normalized from 4.88 million independent clones from a brain library. RNA was made from brain tissue removed from a 26-year-old Caucasian male during cranioplasty and excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated a grade 4 oligoastrocytoma in the right fronto-parietal part of the brain.
43	HIPONON02	This normalized hippocampus library was constructed from 1.13 million independent clones from a hippocampal library. RNA was isolated from the hippocampus tissue of a 72-year-old Caucasian female who died from an intracranial bleed. Patient history included nose cancer, hypertension, and arthritis. The normalization and hybridization conditions were adapted from Soares et al. (PNAS (1994) 91:9928).
44	ADRENOT07	Library was constructed using RNA isolated from adrenal tissue removed from a 61-year-old female during a bilateral adrenalectomy. Patient history included an unspecified disorder of the adrenal glands.
45	GBLANOT01	Library was constructed using RNA isolated from diseased gallbladder tissue removed from a 53-year-old Caucasian female during a cholecystectomy. Pathology indicated mild chronic cholecystitis and cholelithiasis with approximately 150 mixed gallstones. Family history included benign hypertension.
46	KERANOT02	Library was constructed using RNA isolated from epidermal breast keratinocytes (NHEK). NHEK (Clontech #CC-2501) is human breast keratinocyte cell line derived from a 30-year-old black female during breast-reduction surgery.
47	BLADTUT08	Library was constructed using RNA isolated from bladder tumor tissue removed from a 72-year-old Caucasian male during a radical cystectomy and prostatectomy. Pathology indicated an invasive grade 3 (of 3) transitional cell carcinoma in the right bladder base. Patient history included pure hypercholesterolemia and tobacco abuse. Family history included myocardial infarction, cerebrovascular disease, brain cancer, and myocardial infarction.
48	BRSTTUT13	Library was constructed using RNA isolated from breast tumor tissue removed from the right breast of a 46-year-old Caucasian female during a unilateral extended simple mastectomy with breast reconstruction. Pathology indicated an invasive grade 3 adenocarcinoma, ductal type with apocrine features and greater than 50% intraductal component. Patient history included breast cancer.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
49	LUNGNOT31	Library was constructed using RNA isolated from right middle lobe lung tissue removed from a 63-year-old Caucasian male. Pathology for the associated tumor indicated grade 3 adenocarcinoma. Patient history included an abdominal aortic aneurysm, cardiac dysrhythmia, atherosclerotic coronary artery disease, hiatal hernia, chronic sinusitis, and lupus. Family history included acute myocardial infarction and atherosclerotic coronary artery disease.
50	NOSEDIT02	The library was constructed using RNA isolated from nasal polyp tissue.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-25,

b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-25,

c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25, and

10 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-25.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-25.

15 3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide of claim 3 selected from the group consisting of SEQ ID NO:26-50.

20 5. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

6. A cell transformed with a recombinant polynucleotide of claim 5.

25 7. A transgenic organism comprising a recombinant polynucleotide of claim 5.

8. A method for producing a polypeptide of claim 1, the method comprising:

30 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

35 9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:26-50,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

- a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

16. A method for treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

- b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

5

19. A method for treating a disease or condition associated with decreased expression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

10

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

15

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

20

22. A method for treating a disease or condition associated with overexpression of functional EXMAD, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

25

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
 BANDMAN, Olga
 HILLMAN, Jennifer L.
 TANG, Y. Tom
 LAL, Preeti
 YUE, Henry
 BAUGHN, Mariah R.
 LU, Dyung Aina M.
 AZIMZAI, Yalda

<120> EXTRACELLULAR MATRIX AND ADHESION-ASSOCIATED PROTEINS

<130> PF-0693 PCT

<140> To Be Assigned
 <141> Herewith

<150> 60/133,643; 60/150,409
 <151> 1999-05-11; 1999-08-23

<160> 50
 <170> PERL Program

<210> 1
 <211> 309
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 398269CD1

<400> 1
 Met Val Phe Pro Ala Lys Arg Phe Cys Leu Val Pro Ser Met Glu
 1 5 10 15
 Gly Val Arg Trp Ala Phe Ser Cys Gly Thr Trp Leu Pro Ser Arg
 20 25 30
 Ala Glu Trp Leu Leu Ala Val Arg Ser Ile Gln Pro Glu Glu Lys
 35 40 45
 Glu Arg Ile Gly Gln Phe Val Phe Ala Arg Asp Ala Lys Ala Ala
 50 55 60
 Met Ala Gly Arg Leu Met Ile Arg Lys Leu Val Ala Glu Lys Leu
 65 70 75
 Asn Ile Pro Trp Asn His Ile Arg Leu Gln Arg Thr Ala Lys Gly
 80 85 90
 Lys Pro Val Leu Ala Lys Asp Ser Ser Asn Pro Tyr Pro Asn Phe
 95 100 105
 Asn Phe Asn Ile Ser His Gln Gly Asp Tyr Ala Val Leu Ala Ala
 110 115 120
 Glu Pro Glu Leu Gln Val Gly Ile Asp Ile Met Lys Thr Ser Phe
 125 130 135
 Pro Gly Arg Gly Ser Ile Pro Glu Phe Phe His Ile Met Lys Arg
 140 145 150
 Lys Phe Thr Asn Lys Glu Trp Glu Thr Ile Arg Ser Phe Lys Asp
 155 160 165
 Glu Trp Thr Gln Leu Asp Met Phe Tyr Arg Asn Trp Ala Leu Lys
 170 175 180
 Glu Ser Phe Ile Lys Ala Ile Gly Val Gly Leu Gly Phe Glu Leu
 185 190 195
 Gln Arg Leu Glu Phe Asp Leu Ser Pro Leu Asn Leu Asp Ile Gly
 200 205 210
 Gln Val Tyr Lys Glu Thr Arg Leu Phe Leu Asp Gly Glu Glu Glu
 215 220 225
 Lys Glu Trp Ala Phe Glu Glu Ser Lys Ile Asp Glu His His Phe

Val Ala Val Ala	230	Arg Lys Pro Asp	235	Gly Ser Arg His Gln	240
	245		250		255
Val Pro Ser Gln	260	Asp Asp Ser Lys Pro	265	Thr Gln Arg Gln Phe	270
	275		280		285
Ile Leu Asn Phe	290	Asn Asp Leu Met Ser	295	Ser Ala Val Pro Met	300
	305				
Pro Glu Asp Pro		Ser Phe Trp Asp Cys		Phe Cys Phe Thr Glu	
Ile Pro Ile Arg		Asn Gly Thr Lys Ser			

<210> 2

<211> 554

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1258888CD1

<400> 2

Met Pro Leu Pro Trp	1	Ser Leu Ala Leu Pro	10	Leu Leu Leu Ser Trp	15
	5		10		15
Val Ala Gly Gly Phe	20	Gly Asn Ala Ala Ser	25	Ala Arg His His Gly	30
	35		40		45
Leu Leu Ala Ser Ala	50	Arg Gln Pro Gly Val	55	Cys His Tyr Gly Thr	60
	65		70		75
Lys Leu Ala Cys Cys	80	Tyr Gly Trp Arg Arg	85	Asn Ser Lys Gly Val	90
	95		100		105
Cys Glu Ala Thr Cys	110	Glu Pro Gly Cys Lys	115	Phe Gly Glu Cys Val	120
	125		130		135
Gly Pro Asn Lys Cys	140	Arg Cys Phe Pro Gly	145	Tyr Thr Gly Lys Thr	150
	155		160		165
Cys Ser Gln Asp Val	170	Asn Glu Cys Gly Met	175	Lys Pro Arg Pro Cys	180
	185		190		195
Gln His Arg Cys Val	200	Asn Thr His Gly Ser	205	Tyr Lys Cys Phe Cys	210
	215		220		225
Leu Ser Gly His Met	230	Leu Met Pro Asp Ala	235	Thr Cys Val Asn Ser	240
	245		250		255
Arg Thr Cys Ala Met	260	Ile Asn Cys Gln Tyr	265	Ser Cys Glu Asp Thr	270
	275		280		285
Glu Glu Gly Pro Gln	290	Cys Leu Cys Pro Ser	295	Ser Ser Gly Leu Arg Leu	300
	305		310		315
Ala Pro Asn Gly Arg	320	Asp Cys Leu Asp Ile	325	Asp Glu Cys Ala Ser	330
	335		340		345
Gly Lys Val Ile Cys		Pro Tyr Asn Arg Arg		Cys Val Asn Thr Phe	
Gly Ser Tyr Tyr Cys		Lys Cys His Ile Gly		Phe Glu Leu Gln Tyr	
Ile Ser Gly Arg Tyr		Asp Cys Ile Asp Ile		Asn Glu Cys Thr Met	
Asp Ser His Thr Cys		Ser His His Ala Asn		Cys Phe Asn Thr Gln	
Gly Ser Phe Lys Cys		Lys Cys Lys Gln Gly		Tyr Lys Gly Asn Gly	
Leu Arg Cys Ser Ala		Ile Pro Glu Asn Ser		Val Lys Glu Val Leu	
Arg Ala Pro Gly Thr		Ile Lys Asp Arg Ile		Lys Lys Leu Leu Ala	
His Lys Asn Ser Met		Lys Lys Lys Ala Lys		Ile Lys Asn Val Thr	
Pro Glu Pro Thr Arg		Thr Pro Thr Pro Lys		Val Asn Leu Gln Pro	
Phe Asn Tyr Glu Glu		Ile Val Ser Arg Gly		Gly Asn Ser His Gly	
Gly Lys Lys Gly Asn		Glu Glu Lys Met Lys		Glu Gly Leu Glu Asp	

Glu	Lys	Arg	Glu	Glu	Lys	Ala	Leu	Lys	Asn	Asp	Ile	Glu	Glu	Arg
				350					355					360
Ser	Leu	Arg	Gly	Asp	Val	Phe	Phe	Pro	Lys	Val	Asn	Glu	Ala	Gly
				365					370					375
Glu	Phe	Gly	Leu	Ile	Leu	Val	Gln	Arg	Lys	Ala	Leu	Thr	Ser	Lys
				380					385					390
Leu	Glu	His	Lys	Ala	Asp	Leu	Asn	Ile	Ser	Val	Asp	Cys	Ser	Phe
				395					400					405
Asn	His	Gly	Ile	Cys	Asp	Trp	Lys	Gln	Asp	Arg	Glu	Asp	Asp	Phe
				410					415					420
Asp	Trp	Asn	Pro	Ala	Asp	Arg	Asp	Asn	Ala	Ile	Gly	Phe	Tyr	Met
				425					430					435
Ala	Val	Pro	Ala	Leu	Ala	Gly	His	Lys	Lys	Asp	Ile	Gly	Arg	Leu
				440					445					450
Lys	Leu	Leu	Leu	Pro	Asp	Leu	Gln	Pro	Gln	Ser	Asn	Phe	Cys	Leu
				455					460					465
Leu	Phe	Asp	Tyr	Arg	Leu	Ala	Gly	Asp	Lys	Val	Gly	Lys	Leu	Arg
				470					475					480
Val	Phe	Val	Lys	Asn	Ser	Asn	Asn	Ala	Leu	Ala	Trp	Glu	Lys	Thr
				485					490					495
Thr	Ser	Glu	Asp	Glu	Lys	Trp	Lys	Thr	Gly	Lys	Ile	Gln	Leu	Tyr
				500					505					510
Gln	Gly	Thr	Asp	Ala	Thr	Lys	Ser	Ile	Ile	Phe	Glu	Ala	Glu	Arg
				515					520					525
Gly	Lys	Gly	Lys	Thr	Gly	Glu	Ile	Ala	Val	Asp	Gly	Val	Leu	Leu
				530					535					540
Val	Ser	Gly	Leu	Cys	Pro	Asp	Ser	Leu	Leu	Ser	Val	Asp	Asp	
				545					550					

<210> 3

<211> 482

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1375891CD1

<400> 3

Met	Gly	Cys	Leu	Trp	Gly	Leu	Ala	Leu	Pro	Leu	Phe	Phe	Phe	Cys
1				5					10					15
Trp	Glu	Val	Gly	Val	Ser	Gly	Ser	Ser	Ala	Gly	Pro	Ser	Thr	Arg
				20					25					30
Arg	Ala	Asp	Thr	Ala	Met	Thr	Thr	Asp	Asp	Thr	Glu	Val	Pro	Ala
				35					40					45
Met	Thr	Leu	Ala	Pro	Gly	His	Ala	Ala	Leu	Glu	Thr	Gln	Thr	Leu
				50					55					60
Ser	Ala	Glu	Thr	Ser	Ser	Arg	Ala	Ser	Thr	Pro	Ala	Gly	Pro	Ile
				65					70					75
Pro	Glu	Ala	Glu	Thr	Arg	Gly	Ala	Lys	Arg	Ile	Ser	Pro	Ala	Arg
				80					85					90
Glu	Thr	Arg	Ser	Phe	Thr	Lys	Thr	Ser	Pro	Asn	Phe	Met	Val	Leu
				95					100					105
Ile	Ala	Thr	Ser	Val	Glu	Thr	Ser	Ala	Ala	Ser	Gly	Ser	Pro	Glu
				110					115					120
Gly	Ala	Gly	Met	Thr	Thr	Val	Gln	Thr	Ile	Thr	Gly	Ser	Asp	Pro
				125					130					135
Glu	Glu	Ala	Ile	Phe	Asp	Thr	Leu	Cys	Thr	Asp	Asp	Ser	Ser	Glu
				140					145					150
Glu	Ala	Lys	Thr	Leu	Thr	Met	Asp	Ile	Leu	Thr	Leu	Ala	His	Thr
				155					160					165
Ser	Thr	Glu	Ala	Lys	Gly	Leu	Ser	Ser	Glu	Ser	Ser	Ala	Ser	Ser
				170					175					180
Asp	Gly	Pro	His	Pro	Val	Ile	Thr	Pro	Ser	Arg	Ala	Ser	Glu	Ser
				185					190					195
Ser	Ala	Ser	Ser	Asp	Gly	Pro	His	Pro	Val	Ile	Thr	Pro	Ser	Arg
				200					205					210
Ala	Ser	Glu	Ser	Ser	Ala	Ser	Ser	Asp	Gly	Pro	His	Pro	Val	Ile

Thr	Pro	Ser	Trp	215	Ser	Pro	Gly	Ser	Asp	220	Thr	Leu	Leu	Ala	225	Glu
Ala	Leu	Val	Thr	230	Val	Thr	Asn	Ile	Glu	235	Val	Ile	Asn	Cys	240	Ile
Thr	Glu	Ile	Glu	245	Thr	Thr	Ser	Ser	Ile	250	Pro	Gly	Ala	Ser	255	Asp
Ile	Asp	Leu	Ile	260	Pro	Thr	Glu	Gly	Val	265	Lys	Ala	Ser	Ser	270	Ser
Asp	Pro	Pro	Ala	275	Leu	Pro	Asp	Ser	Thr	280	Glu	Ala	Lys	Pro	285	Ile
Thr	Glu	Val	Thr	290	Ala	Ser	Ala	Glu	Thr	295	Leu	Ser	Thr	Ala	300	Thr
Thr	Glu	Ser	Ala	305	Ala	Pro	His	Ala	Thr	310	Val	Gly	Thr	Pro	315	Pro
Thr	Asn	Ser	Ala	320	Thr	Glu	Arg	Glu	Val	325	Ala	Pro	Gly	Ala	330	Thr
Thr	Leu	Ser	Gly	335	Ala	Leu	Val	Thr	Val	340	Ser	Arg	Asn	Pro	345	Glu
Glu	Thr	Ser	Ala	350	Leu	Ser	Val	Glu	Thr	355	Pro	Ser	Tyr	Val	360	Val
Ser	Gly	Ala	Ala	365	Pro	Val	Ser	Ile	Glu	370	Ala	Gly	Ser	Ala	375	Gly
Lys	Thr	Thr	Ser	380	Phe	Ala	Gly	Ser	Ser	385	Ala	Ser	Ser	Tyr	390	Pro
Ser	Glu	Ala	Ala	395	Leu	Lys	Asn	Phe	Thr	400	Pro	Ser	Glu	Thr	405	Thr
Met	Asp	Ile	Ala	410	Thr	Lys	Gly	Pro	Phe	415	Pro	Thr	Ser	Arg	420	Pro
Leu	Pro	Ser	Val	425	Pro	Pro	Thr	Thr	Thr	430	Asn	Ser	Ser	Arg	435	Thr
Asn	Ser	Thr	Leu	440	Ala	Lys	Ile	Thr	Thr	445	Ser	Ala	Lys	Thr	450	Met
Lys	Pro	Gln	Gln	455	Pro	Arg	Pro	Arg	Leu	460	Pro	Gly	Arg	Gly	465	Pro
Gln	Thr			470						475					480	

<210> 4

<211> 735

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1524355CD1

<400> 4

Met	Ala	Ala	Gly	Gly	Ala	Val	Ala	Ala	Ala	Pro	Glu	Cys	Arg	Leu		
1				5					10					15		
Leu	Pro	Tyr	Ala	Leu	His	Lys	Trp	Ser	Ser	Phe	Ser	Ser	Thr	Tyr		
				20					25					30		
Leu	Pro	Glu	Asn	Ile	Leu	Val	Asp	Lys	Pro	Asn	Asp	Gln	Ser	Ser		
				35					40					45		
Arg	Trp	Ser	Ser	Glu	Ser	Asn	Tyr	Pro	Pro	Gln	Tyr	Leu	Ile	Leu		
				50					55					60		
Lys	Leu	Glu	Arg	Pro	Ala	Ile	Val	Gln	Asn	Ile	Thr	Phe	Gly	Lys		
				65					70					75		
Tyr	Glu	Lys	Thr	His	Val	Cys	Asn	Leu	Lys	Lys	Phe	Lys	Val	Phe		
				80					85					90		
Gly	Gly	Met	Asn	Glu	Glu	Asn	Met	Thr	Glu	Leu	Leu	Ser	Ser	Gly		
				95					100					105		
Leu	Lys	Asn	Asp	Tyr	Asn	Lys	Glu	Thr	Phe	Thr	Leu	Lys	His	Lys		
				110					115					120		
Ile	Asp	Glu	Gln	Met	Phe	Pro	Cys	Arg	Phe	Ile	Lys	Ile	Val	Pro		
				125					130					135		
Leu	Leu	Ser	Trp	Gly	Pro	Ser	Phe	Asn	Phe	Ser	Ile	Trp	Tyr	Val		
				140					145					150		

Glu	Leu	Ser	Gly	Ile	Asp	Asp	Pro	Asp	Ile	Val	Gln	Pro	Cys	Leu
				155					160					165
Asn	Trp	Tyr	Ser	Lys	Tyr	Arg	Glu	Gln	Glu	Ala	Ile	Arg	Leu	Cys
				170					175					180
Leu	Lys	His	Phe	Arg	Gln	His	Asn	Tyr	Thr	Glu	Ala	Phe	Glu	Ser
				185					190					195
Leu	Gln	Lys	Lys	Thr	Lys	Ile	Ala	Leu	Glu	His	Pro	Met	Leu	Thr
				200					205					210
Asp	Ile	His	Asp	Lys	Leu	Val	Leu	Lys	Gly	Asp	Phe	Asp	Ala	Cys
				215					220					225
Glu	Glu	Leu	Ile	Glu	Lys	Ala	Val	Asn	Asp	Gly	Leu	Phe	Asn	Gln
				230					235					240
Tyr	Ile	Ser	Gln	Gln	Glu	Tyr	Lys	Pro	Arg	Trp	Ser	Gln	Ile	Ile
				245					250					255
Pro	Lys	Ser	Thr	Lys	Gly	Asp	Gly	Glu	Asp	Asn	Arg	Pro	Gly	Met
				260					265					270
Arg	Gly	Gly	His	Gln	Met	Val	Ile	Asp	Val	Gln	Thr	Glu	Thr	Val
				275					280					285
Tyr	Leu	Phe	Gly	Gly	Trp	Asp	Gly	Thr	Gln	Asp	Leu	Ala	Asp	Phe
				290					295					300
Trp	Ala	Tyr	Ser	Val	Lys	Glu	Asn	Gln	Trp	Thr	Cys	Ile	Ser	Arg
				305					310					315
Asp	Thr	Glu	Lys	Glu	Asn	Gly	Pro	Ser	Ala	Arg	Ser	Cys	His	Lys
				320					325					330
Met	Cys	Ile	Asp	Ile	Gln	Arg	Arg	Gln	Ile	Tyr	Thr	Leu	Gly	Arg
				335					340					345
Tyr	Leu	Asp	Ser	Ser	Val	Arg	Asn	Ser	Lys	Ser	Leu	Lys	Ser	Asp
				350					355					360
Phe	Tyr	Arg	Tyr	Asp	Ile	Asp	Thr	Asn	Thr	Trp	Met	Leu	Leu	Ser
				365					370					375
Glu	Asp	Thr	Ala	Ala	Asp	Gly	Gly	Pro	Lys	Leu	Val	Phe	Asp	His
				380					385					390
Gln	Met	Cys	Met	Asp	Ser	Glu	Lys	His	Met	Ile	Tyr	Thr	Phe	Gly
				395					400					405
Gly	Arg	Ile	Leu	Thr	Cys	Asn	Gly	Ser	Val	Asp	Asp	Ser	Arg	Ala
				410					415					420
Ser	Glu	Pro	Gln	Phe	Ser	Gly	Leu	Phe	Ala	Phe	Asn	Cys	Gln	Cys
				425					430					435
Gln	Thr	Trp	Lys	Leu	Leu	Arg	Glu	Asp	Ser	Cys	Asn	Ala	Gly	Pro
				440					445					450
Glu	Asp	Ile	Gln	Ser	Arg	Ile	Gly	His	Cys	Met	Leu	Phe	His	Ser
				455					460					465
Lys	Asn	Arg	Cys	Leu	Tyr	Val	Phe	Gly	Gly	Gln	Arg	Ser	Lys	Thr
				470					475					480
Tyr	Leu	Asn	Asp	Phe	Phe	Ser	Tyr	Asp	Val	Asp	Ser	Asp	His	Val
				485					490					495
Asp	Ile	Ile	Ser	Asp	Gly	Thr	Lys	Lys	Asp	Ser	Gly	Met	Val	Pro
				500					505					510
Met	Thr	Gly	Phe	Thr	Gln	Arg	Ala	Thr	Ile	Asp	Pro	Glu	Leu	Asn
				515					520					525
Glu	Ile	His	Val	Leu	Ser	Gly	Leu	Ser	Lys	Asp	Lys	Glu	Lys	Arg
				530					535					540
Glu	Glu	Asn	Val	Arg	Asn	Ser	Phe	Trp	Ile	Tyr	Asp	Ile	Val	Arg
				545					550					555
Asn	Ser	Trp	Ser	Cys	Val	Tyr	Lys	Asn	Asp	Gln	Ala	Ala	Lys	Asp
				560					565					570
Asn	Pro	Thr	Lys	Ser	Leu	Gln	Glu	Glu	Glu	Pro	Cys	Pro	Arg	Phe
				575					580					585
Ala	His	Gln	Leu	Val	Tyr	Asp	Glu	Leu	His	Lys	Val	His	Tyr	Leu
				590					595					600
Phe	Gly	Gly	Asn	Pro	Gly	Lys	Ser	Cys	Ser	Pro	Lys	Met	Arg	Leu
				605					610					615
Asp	Asp	Phe	Trp	Ser	Leu	Lys	Leu	Cys	Arg	Pro	Ser	Lys	Asp	Tyr
				620					625					630
Leu	Leu	Arg	His	Cys	Lys	Tyr	Leu	Ile	Arg	Lys	His	Arg	Phe	Glu
				635					640					645
Glu	Lys	Ala	Gln	Val	Asp	Pro	Leu	Ser	Ala	Leu	Lys	Tyr	Leu	Gln

Asn Asp Leu Tyr	650	Thr Val Asp His	655	Ser Asp Pro Glu Glu	660
	665		670		675
Lys Glu Phe Gln	680	Leu Leu Ala Ser Ala	685	Leu Phe Lys Ser Gly	690
	695		700		705
Asp Phe Thr Ala	710	Leu Gly Phe Ser Asp	715	Val Asp His Thr Tyr	720
	725		730		735
Gln Arg Thr Gln		Leu Phe Asp Thr Leu		Val Asn Phe Phe Pro	
Ser Met Thr Pro		Pro Lys Gly Asn Leu		Val Asp Leu Ile Thr	

<210> 5

<211> 424

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1598937CD1

<400> 5

Met Ala Pro Glu Glu	5	Asp Ala Gly Gly	10	Glu Ala Leu Gly Gly	15
1					
Phe Trp Glu Ala Gly	20	Asn Tyr Arg Arg	25	Thr Val Gln Arg Val	30
	35		40		45
Asp Gly His Arg Leu	50	Cys Gly Asp Leu	55	Ser Cys Phe Gln Glu	60
	65		70		75
Arg Ala Arg Ile Glu	80	Lys Ala Tyr Ala	85	Gln Gln Leu Ala Asp	90
	95		100		105
Ala Arg Lys Trp Arg	110	Gly Thr Val Glu	115	Gly Pro Gln Tyr Gly	120
	125		130		135
Thr Leu Glu Lys Ala	140	Trp His Ala Phe	145	Phe Thr Ala Ala Glu	150
	155		160		165
Leu Ser Ala Leu His	170	Leu Glu Val Arg	175	Glu Lys Leu Gln Gly	180
	185		190		195
Asp Ser Glu Arg Val	200	Arg Ala Trp Gln	205	Arg Gly Ala Phe His	210
	215		220		225
Pro Val Leu Gly Gly	230	Phe Arg Glu Ser	235	Arg Ala Ala Glu Asp	240
	245		250		255
Phe Arg Lys Ala Gln	260	Lys Pro Trp Leu	265	Lys Arg Leu Lys Glu	270
	275		280		285
Glu Ala Ser Lys Lys	290	Ser Tyr His Ala	295	Ala Arg Lys Asp Glu	300
	305		310		315
Thr Ala Gln Thr Arg	320	Glu Ser His Ala	325	Lys Ala Asp Ser Ala	330
	335		340		345
Ser Gln Glu Gln Leu		Arg Lys Leu Gln		Glu Arg Val Glu Arg	
Ala Lys Glu Ala Glu		Lys Thr Lys Ala		Gln Tyr Glu Gln Thr	
Ala Glu Leu His Arg		Tyr Thr Pro Arg		Met Glu Asp Met Glu	
Gln Ala Phe Glu Thr		Cys Gln Ala Ala		Glu Arg Gln Arg Leu	
Phe Phe Lys Asp Met		Leu Leu Thr Leu		His Gln His Leu Asp	
Ser Ser Ser Glu Lys		Phe His Glu Leu		His Arg Asp Leu His	
Gly Ile Glu Ala Ala		Ser Asp Glu Glu		Asp Leu Arg Trp Trp	
Ser Thr His Gly Pro		Gly Met Ala Met		Asn Trp Pro Gln Phe	
Glu Trp Ser Leu Asp		Thr Gln Arg Thr		Ile Ser Arg Lys Glu	
Gly Gly Arg Ser Pro		Asp Glu Val Thr		Leu Thr Ser Ile Val	
Thr Arg Asp Gly Thr		Ala Pro Pro Pro		Gln Ser Pro Gly Ser	

Gly Thr Gly Gln	Asp Glu Glu Trp Ser	Asp Glu Glu Ser Pro	Arg
	350	355	360
Lys Ala Ala Thr	Gly Val Arg Val Arg	Ala Leu Tyr Asp Tyr	Ala
	365	370	375
Gly Gln Glu Ala	Asp Glu Leu Ser Phe	Arg Ala Gly Glu Glu	Leu
	380	385	390
Leu Lys Met Ser	Glu Glu Asp Glu Gln	Gly Trp Cys Gln Gly	Gln
	395	400	405
Leu Gln Ser Gly	Arg Ile Gly Leu Tyr	Pro Ala Asn Tyr Val	Glu
	410	415	420
Cys Val Gly Ala			

<210> 6

<211> 420

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1725801CD1

<400> 6

Met Ala Pro Trp	Pro Pro Lys Gly Leu Val	Pro Ala Val Leu Trp
1	5	10
Gly Leu Ser Leu	Phe Leu Asn Leu Pro	Gly Pro Ile Trp Leu Gln
	20	25
Pro Ser Pro Pro	Pro Gln Ser Ser Pro	Pro Pro Gln Pro His Pro
	35	40
Cys His Thr Cys	Arg Gly Leu Val Asp	Ser Phe Asn Lys Gly Leu
	50	55
Glu Arg Thr Ile	Arg Asp Asn Phe Gly	Gly Gly Asn Thr Ala Trp
	65	70
Glu Glu Glu Asn	Leu Ser Lys Tyr Lys	Asp Ser Glu Thr Arg Leu
	80	85
Val Glu Val Leu	Glu Gly Val Cys Ser	Lys Ser Asp Phe Glu Cys
	95	100
His Arg Leu Leu	Glu Leu Ser Glu Glu	Leu Val Glu Ser Trp Trp
	110	115
Phe His Lys Gln	Gln Glu Ala Pro Asp	Leu Phe Gln Trp Leu Cys
	125	130
Ser Asp Ser Leu	Lys Leu Cys Cys Pro	Ala Gly Thr Phe Gly Pro
	140	145
Ser Cys Leu Pro	Cys Pro Gly Gly Thr	Glu Arg Pro Cys Gly Gly
	155	160
Tyr Gly Gln Cys	Glu Gly Glu Gly Thr	Arg Gly Gly Ser Gly His
	170	175
Cys Asp Cys Gln	Ala Gly Tyr Gly Gly	Glu Ala Cys Gly Gln Cys
	185	190
Gly Leu Gly Tyr	Phe Glu Ala Glu Arg	Asn Ala Ser His Leu Val
	200	205
Cys Ser Ala Cys	Phe Gly Pro Cys Ala	Arg Cys Ser Gly Pro Glu
	215	220
Glu Ser Asn Cys	Leu Gln Cys Lys Lys	Gly Trp Ala Leu His His
	230	235
Leu Lys Cys Val	Asp Ile Asp Glu Cys	Gly Thr Glu Gly Ala Asn
	245	250
Cys Gly Ala Asp	Gln Phe Cys Val Asn	Thr Glu Gly Ser Tyr Glu
	260	265
Cys Arg Asp Cys	Ala Lys Ala Cys Leu	Gly Cys Met Gly Ala Gly
	275	280
Pro Gly Arg Cys	Lys Lys Cys Ser Pro	Gly Tyr Gln Gln Val Gly
	290	295
Ser Lys Cys Leu	Asp Val Asp Glu Cys	Glu Thr Glu Val Cys Pro
	305	310
Gly Glu Asn Lys	Gln Cys Glu Asn Thr	Glu Gly Gly Tyr Arg Cys
	320	325
Ile Cys Ala Glu	Gly Tyr Lys Gln Met	Glu Gly Ile Cys Val Lys

Glu	Gln	Ile	Pro	335	Glu	Ser	Ala	Gly	Phe	340	Ser	Glu	Met	Thr	345	Glu
Asp	Glu	Leu	Val	350	Val	Leu	Gln	Gln	Met	355	Phe	Gly	Ile	Ile	360	Ile
Cys	Ala	Leu	Ala	365	Thr	Leu	Ala	Ala	Lys	370	Gly	Asp	Leu	Val	375	Thr
Ala	Ile	Phe	Ile	380	Gly	Ala	Val	Ala	Ala	385	Met	Thr	Gly	Tyr	390	Leu
Ser	Glu	Arg	Ser	395	Asp	Arg	Val	Leu	Glu	400	Gly	Phe	Ile	Lys	405	Arg
				410						415					420	

<210> 7

<211> 795

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1730482CD1

<400> 7

Met	Glu	Lys	Thr	Gln	Ser	Leu	Pro	Thr	Arg	Pro	Pro	Thr	Phe	Pro
1				5					10					15
Pro	Thr	Ile	Pro	Pro	Ala	Lys	Glu	Val	Cys	Lys	Ala	Ala	Lys	Ala
				20					25					30
Asp	Leu	Val	Phe	Met	Val	Asp	Gly	Ser	Trp	Ser	Ile	Gly	Asp	Glu
				35					40					45
Asn	Phe	Asn	Lys	Ile	Ile	Ser	Phe	Leu	Tyr	Ser	Thr	Val	Gly	Ala
				50					55					60
Leu	Asn	Lys	Ile	Gly	Thr	Asp	Gly	Thr	Gln	Val	Ala	Met	Val	Gln
				65					70					75
Phe	Thr	Asp	Asp	Pro	Arg	Thr	Glu	Phe	Lys	Leu	Asn	Ala	Tyr	Lys
				80					85					90
Thr	Lys	Glu	Thr	Leu	Leu	Asp	Ala	Ile	Lys	His	Ile	Ser	Tyr	Lys
				95					100					105
Gly	Gly	Asn	Thr	Lys	Thr	Gly	Lys	Ala	Ile	Lys	Tyr	Val	Arg	Asp
				110					115					120
Thr	Leu	Phe	Thr	Ala	Glu	Ser	Gly	Thr	Arg	Arg	Gly	Ile	Pro	Lys
				125					130					135
Val	Ile	Val	Val	Ile	Thr	Asp	Gly	Arg	Ser	Gln	Asp	Asp	Val	Asn
				140					145					150
Lys	Ile	Ser	Arg	Glu	Met	Gln	Leu	Asp	Gly	Tyr	Ser	Ile	Phe	Ala
				155					160					165
Ile	Gly	Val	Ala	Asp	Ala	Asp	Tyr	Ser	Glu	Leu	Val	Ser	Ile	Gly
				170					175					180
Ser	Lys	Pro	Ser	Ala	Arg	His	Val	Phe	Phe	Val	Asp	Asp	Phe	Asp
				185					190					195
Ala	Phe	Lys	Lys	Ile	Glu	Asp	Glu	Leu	Ile	Thr	Phe	Val	Cys	Glu
				200					205					210
Thr	Ala	Ser	Ala	Thr	Cys	Pro	Val	Val	His	Lys	Asp	Gly	Ile	Asp
				215					220					225
Leu	Ala	Gly	Phe	Lys	Met	Met	Glu	Met	Phe	Gly	Leu	Val	Glu	Lys
				230					235					240
Asp	Phe	Ser	Ser	Val	Glu	Gly	Val	Ser	Met	Glu	Pro	Gly	Thr	Phe
				245					250					255
Asn	Val	Phe	Pro	Cys	Tyr	Gln	Leu	His	Lys	Asp	Ala	Leu	Val	Ser
				260					265					270
Gln	Pro	Thr	Arg	Tyr	Leu	His	Pro	Glu	Gly	Leu	Pro	Ser	Asp	Tyr
				275					280					285
Thr	Ile	Ser	Phe	Leu	Phe	Arg	Ile	Leu	Pro	Asp	Thr	Pro	Gln	Glu
				290					295					300
Pro	Phe	Ala	Leu	Trp	Glu	Ile	Leu	Asn	Lys	Asn	Ser	Asp	Pro	Leu
				305					310					315
Val	Gly	Val	Ile	Leu	Asp	Asn	Gly	Gly	Lys	Thr	Leu	Thr	Tyr	Phe
				320					325					330
Asn	Tyr	Asp	Gln	Ser	Gly	Asp	Phe	Gln	Thr	Val	Thr	Phe	Glu	Gly
				335					340					345

Pro	Glu	Ile	Arg	Lys	Ile	Phe	Tyr	Gly	Ser	Phe	His	Lys	Leu	His
				350					355					360
Ile	Val	Val	Ser	Glu	Thr	Leu	Val	Lys	Val	Val	Ile	Asp	Cys	Lys
				365					370					375
Gln	Val	Gly	Glu	Lys	Ala	Met	Asn	Ala	Ser	Ala	Asn	Ile	Thr	Ser
				380					385					390
Asp	Gly	Val	Glu	Val	Leu	Gly	Lys	Met	Val	Arg	Ser	Arg	Gly	Pro
				395					400					405
Gly	Gly	Asn	Ser	Ala	Pro	Phe	Gln	Leu	Gln	Met	Phe	Asp	Ile	Val
				410					415					420
Cys	Ser	Thr	Ser	Trp	Ala	Asn	Thr	Asp	Lys	Cys	Cys	Glu	Leu	Pro
				425					430					435
Gly	Leu	Arg	Asp	Asp	Glu	Ser	Cys	Pro	Asp	Leu	Pro	His	Ser	Cys
				440					445					450
Ser	Cys	Ser	Glu	Thr	Asn	Glu	Val	Ala	Leu	Gly	Pro	Ala	Gly	Pro
				455					460					465
Pro	Gly	Gly	Pro	Gly	Leu	Arg	Gly	Pro	Lys	Gly	Gln	Gln	Gly	Glu
				470					475					480
Pro	Gly	Pro	Lys	Gly	Pro	Asp	Gly	Pro	Arg	Gly	Glu	Ile	Gly	Leu
				485					490					495
Pro	Gly	Pro	Gln	Gly	Pro	Pro	Gly	Pro	Gln	Gly	Pro	Ser	Gly	Leu
				500					505					510
Ser	Ile	Gln	Gly	Met	Pro	Gly	Met	Pro	Gly	Glu	Lys	Gly	Glu	Lys
				515					520					525
Gly	Asp	Thr	Gly	Leu	Pro	Gly	Pro	Gln	Gly	Ile	Pro	Gly	Gly	Val
				530					535					540
Gly	Ser	Pro	Gly	Arg	Asp	Gly	Ser	Pro	Gly	Gln	Arg	Gly	Leu	Pro
				545					550					555
Gly	Lys	Asp	Gly	Ser	Ser	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro	Ile
				560					565					570
Gly	Ile	Pro	Gly	Thr	Pro	Gly	Val	Pro	Gly	Ile	Thr	Gly	Ser	Met
				575					580					585
Gly	Pro	Gln	Gly	Ala	Leu	Gly	Pro	Pro	Gly	Val	Pro	Gly	Ala	Lys
				590					595					600
Gly	Glu	Arg	Gly	Glu	Arg	Gly	Asp	Leu	Gln	Ser	Gln	Ala	Met	Val
				605					610					615
Arg	Ser	Val	Ala	Arg	Gln	Val	Cys	Glu	Gln	Leu	Ile	Gln	Ser	His
				620					625					630
Met	Ala	Arg	Tyr	Thr	Ala	Ile	Leu	Asn	Gln	Ile	Pro	Ser	His	Ser
				635					640					645
Ser	Ser	Ile	Arg	Thr	Val	Gln	Gly	Pro	Pro	Gly	Glu	Pro	Gly	Arg
				650					655					660
Pro	Gly	Ser	Pro	Gly	Ala	Pro	Gly	Glu	Gln	Gly	Pro	Pro	Gly	Thr
				665					670					675
Pro	Gly	Phe	Pro	Gly	Asn	Ala	Gly	Val	Pro	Gly	Thr	Pro	Gly	Glu
				680					685					690
Arg	Gly	Leu	Thr	Gly	Ile	Lys	Gly	Glu	Lys	Gly	Asn	Pro	Gly	Val
				695					700					705
Gly	Thr	Gln	Gly	Pro	Arg	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Pro	Ser
				710					715					720
Gly	Glu	Ser	Arg	Pro	Gly	Ser	Pro	Gly	Pro	Pro	Gly	Ser	Pro	Gly
				725					730					735
Pro	Arg	Gly	Pro	Pro	Gly	His	Leu	Gly	Val	Pro	Gly	Pro	Gln	Gly
				740					745					750
Pro	Ser	Gly	Gln	Pro	Gly	Tyr	Cys	Asp	Pro	Ser	Ser	Cys	Ser	Ala
				755					760					765
Tyr	Gly	Val	Arg	Ala	Pro	His	Pro	Asp	Gln	Pro	Glu	Phe	Thr	Pro
				770					775					780
Val	Gln	Asp	Glu	Leu	Glu	Ala	Met	Glu	Leu	Trp	Gly	Pro	Gly	Val
				785					790					795

<210> 8

<211> 306

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1810058CD1

<400> 8

```

Met Arg Ile Trp Trp Leu Leu Leu Ala Ile Glu Ile Cys Thr Gly
 1          5          10          15
Asn Ile Asn Ser Gln Asp Thr Cys Arg Gln Gly His Pro Gly Ile
 20          25          30
Pro Gly Asn Pro Gly His Asn Gly Leu Pro Gly Arg Asp Gly Arg
 35          40          45
Asp Gly Ala Lys Gly Asp Lys Gly Asp Ala Gly Glu Pro Gly Arg
 50          55          60
Pro Gly Ser Pro Gly Lys Asp Gly Thr Ser Gly Glu Lys Gly Glu
 65          70          75
Arg Gly Ala Asp Gly Lys Val Glu Ala Lys Gly Ile Lys Gly Asp
 80          85          90
Gln Gly Ser Arg Gly Ser Pro Gly Lys His Gly Pro Lys Gly Leu
 95          100          105
Ala Gly Pro Met Gly Glu Lys Gly Leu Arg Gly Glu Thr Gly Pro
110          115          120
Gln Gly Gln Lys Gly Asn Lys Gly Asp Val Gly Pro Thr Gly Pro
125          130          135
Glu Gly Pro Arg Gly Asn Ile Gly Pro Leu Gly Pro Thr Gly Leu
140          145          150
Pro Gly Pro Met Gly Pro Ile Gly Lys Pro Gly Pro Lys Gly Glu
155          160          165
Ala Gly Pro Thr Gly Pro Gln Gly Glu Pro Gly Val Arg Gly Ile
170          175          180
Arg Gly Trp Lys Gly Asp Arg Gly Glu Lys Gly Lys Ile Gly Glu
185          190          195
Thr Leu Val Leu Pro Lys Ser Ala Phe Thr Val Gly Leu Thr Val
200          205          210
Leu Ser Lys Phe Pro Ser Ser Asp Val Pro Ile Lys Phe Asp Lys
215          220          225
Ile His Ile Thr Val Phe Ser Arg Asn Val Gln Val Ser Leu Val
230          235          240
Lys Asn Gly Val Lys Ile Leu His Thr Arg Asp Ala Tyr Val Ser
245          250          255
Ser Glu Asp Gln Ala Ser Gly Ser Ile Val Leu Gln Leu Lys Leu
260          265          270
Gly Asp Glu Met Trp Leu Gln Val Thr Gly Gly Glu Arg Phe Asn
275          280          285
Gly Leu Phe Ala Asp Glu Asp Asp Asp Thr Thr Phe Thr Gly Phe
290          295          300
Leu Leu Phe Ser Ser Gln
305

```

<210> 9

<211> 338

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2040679CD1

<400> 9

```

Met Tyr Val Leu Ser Pro Val Glu Phe Ile Ile Leu Gln Leu Leu
 1          5          10          15
Phe Ile Gln Ala Ile Ser Ser Ser Leu Lys Gly Phe Leu Ser Ala
 20          25          30
Met Arg Leu Ala His Arg Gly Cys Asn Val Asp Thr Pro Val Ser
 35          40          45
Thr Leu Thr Pro Val Lys Thr Ser Glu Phe Glu Asn Phe Lys Thr
 50          55          60
Lys Met Val Ile Thr Ser Lys Lys Asp Tyr Pro Leu Ser Lys Asn
 65          70          75
Phe Pro Tyr Ser Leu Glu His Leu Gln Thr Ser Tyr Cys Gly Leu
 80          85          90

```

Val	Arg	Val	Asp	Met	Arg	Met	Leu	Cys	Leu	Lys	Ser	Leu	Arg	Lys
				95					100					105
Leu	Asp	Leu	Ser	His	Asn	His	Ile	Lys	Lys	Leu	Pro	Ala	Thr	Ile
				110					115					120
Gly	Asp	Leu	Ile	His	Leu	Gln	Glu	Leu	Asn	Leu	Asn	Asp	Asn	His
				125					130					135
Leu	Glu	Ser	Phe	Ser	Val	Ala	Leu	Cys	His	Ser	Thr	Leu	Gln	Lys
				140					145					150
Ser	Leu	Arg	Ser	Leu	Asp	Leu	Ser	Lys	Asn	Lys	Ile	Lys	Ala	Leu
				155					160					165
Pro	Val	Gln	Phe	Cys	Gln	Leu	Gln	Glu	Leu	Lys	Asn	Leu	Lys	Leu
				170					175					180
Asp	Asp	Asn	Glu	Leu	Ile	Gln	Phe	Pro	Cys	Lys	Ile	Gly	Gln	Leu
				185					190					195
Ile	Asn	Leu	Arg	Phe	Leu	Ser	Ala	Ala	Arg	Asn	Lys	Leu	Pro	Phe
				200					205					210
Leu	Pro	Ser	Glu	Phe	Arg	Asn	Leu	Ser	Leu	Glu	Tyr	Leu	Asp	Leu
				215					220					225
Phe	Gly	Asn	Thr	Phe	Glu	Gln	Pro	Lys	Val	Leu	Pro	Val	Ile	Lys
				230					235					240
Leu	Gln	Ala	Pro	Leu	Thr	Leu	Leu	Glu	Ser	Ser	Ala	Arg	Thr	Ile
				245					250					255
Leu	His	Asn	Arg	Ile	Pro	Tyr	Gly	Ser	His	Ile	Ile	Pro	Phe	His
				260					265					270
Leu	Cys	Gln	Asp	Leu	Asp	Thr	Ala	Lys	Ile	Cys	Val	Cys	Gly	Arg
				275					280					285
Phe	Cys	Leu	Asn	Ser	Phe	Ile	Gln	Gly	Thr	Thr	Thr	Met	Asn	Leu
				290					295					300
His	Ser	Val	Ala	His	Thr	Val	Val	Leu	Val	Asp	Asn	Leu	Gly	Gly
				305					310					315
Thr	Glu	Ala	Pro	Ile	Ile	Ser	Tyr	Phe	Cys	Ser	Leu	Gly	Cys	Tyr
				320					325					330
Val	Asn	Ser	Ser	Asp	Met	Leu	Lys							
				335										

<210> 10

<211> 164

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2960051CD1

<400> 10

Met	Lys	Ile	Ala	Val	Leu	Phe	Cys	Phe	Phe	Leu	Leu	Ile	Ile	Phe
1				5					10					15
Gln	Thr	Asp	Phe	Gly	Lys	Asn	Glu	Glu	Ile	Pro	Arg	Lys	Gln	Arg
				20					25					30
Arg	Lys	Ile	Tyr	His	Arg	Arg	Leu	Arg	Lys	Ser	Ser	Thr	Ser	His
				35					40					45
Lys	His	Arg	Ser	Asn	Arg	Gln	Leu	Gly	Ile	Pro	Gln	Thr	Thr	Val
				50					55					60
Phe	Thr	Pro	Val	Ala	Arg	Leu	Pro	Ile	Val	Asn	Phe	Asp	Tyr	Ser
				65					70					75
Met	Glu	Glu	Lys	Phe	Glu	Ser	Phe	Ser	Ser	Phe	Pro	Gly	Val	Glu
				80					85					90
Ser	Ser	Tyr	Asn	Val	Leu	Pro	Gly	Lys	Lys	Gly	His	Cys	Leu	Val
				95					100					105
Lys	Gly	Ile	Thr	Met	Tyr	Asn	Lys	Ala	Val	Trp	Ser	Pro	Glu	Pro
				110					115					120
Cys	Thr	Thr	Cys	Leu	Cys	Ser	Asp	Gly	Arg	Val	Leu	Cys	Asp	Glu
				125					130					135
Thr	Met	Cys	His	Pro	Gln	Arg	Cys	Pro	Gln	Thr	Val	Ile	Pro	Glu
				140					145					150
Gly	Glu	Cys	Cys	Pro	Val	Cys	Ser	Ala	Thr	Gly	Thr	Glu	Ile	
				155					160					

<210> 11

<211> 327

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3117318CD1

<400> 11

```

Met Arg Ala Leu Pro Gly Leu Leu Glu Ala Arg Ala Arg Thr Pro
 1      5      10
Arg Leu Leu Leu Leu Gln Cys Leu Leu Ala Ala Ala Arg Pro Ser
      20      25      30
Ser Ala Asp Gly Ser Ala Pro Asp Ser Ala Phe Thr Ser Pro Pro
      35      40      45
Leu Arg Glu Glu Ile Met Ala Asn Asn Phe Ser Leu Glu Ser His
      50      55      60
Asn Ile Ser Leu Thr Glu His Ser Ser Met Pro Val Glu Lys Asn
      65      70      75
Ile Thr Leu Glu Arg Pro Ser Asn Val Asn Leu Thr Cys Gln Phe
      80      85      90
Thr Thr Ser Gly Asp Leu Asn Ala Val Asn Val Thr Trp Lys Lys
      95      100      105
Asp Gly Glu Gln Leu Glu Asn Asn Tyr Leu Val Ser Ala Thr Gly
      110      115      120
Ser Thr Leu Tyr Thr Gln Tyr Arg Phe Thr Ile Ile Asn Ser Lys
      125      130      135
Gln Met Gly Ser Tyr Ser Cys Phe Phe Arg Glu Glu Lys Glu Gln
      140      145      150
Arg Gly Thr Phe Asn Phe Lys Val Pro Glu Leu His Gly Lys Asn
      155      160      165
Lys Pro Leu Ile Ser Tyr Val Gly Asp Ser Thr Val Leu Thr Cys
      170      175      180
Lys Cys Gln Asn Cys Phe Pro Leu Asn Trp Thr Trp Tyr Ser Ser
      185      190      195
Asn Gly Ser Val Lys Val Pro Val Gly Val Gln Met Asn Lys Tyr
      200      205      210
Val Ile Asn Gly Thr Tyr Ala Asn Glu Thr Lys Leu Lys Ile Thr
      215      220      225
Gln Leu Leu Glu Glu Asp Gly Glu Ser Tyr Trp Cys Arg Ala Leu
      230      235      240
Phe Gln Leu Gly Glu Ser Glu Glu His Ile Glu Leu Val Val Leu
      245      250      255
Ser Tyr Leu Val Pro Leu Lys Pro Phe Leu Val Ile Val Ala Glu
      260      265      270
Val Ile Leu Leu Val Ala Thr Ile Leu Leu Cys Glu Lys Tyr Thr
      275      280      285
Gln Lys Lys Lys Lys His Ser Asp Glu Gly Lys Glu Phe Glu Gln
      290      295      300
Ile Glu Gln Leu Lys Ser Asp Asp Ser Asn Gly Ile Glu Asn Asn
      305      310      315
Val Pro Arg His Arg Lys Asn Glu Ser Leu Gly Gln
      320      325

```

<210> 12

<211> 716

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3486992CD1

<400> 12

```

Met Ala Arg Met Ser Phe Val Ile Ala Ala Cys Gln Leu Val Leu
 1      5      10
Gly Leu Leu Met Thr Ser Leu Thr Glu Ser Ser Ile Gln Asn Ser
      20      25      30

```

Glu	Cys	Pro	Gln	Leu	Cys	Val	Cys	Glu	Ile	Arg	Pro	Trp	Phe	Thr	
				35					40						45
Pro	Gln	Ser	Thr	Tyr	Arg	Glu	Ala	Thr	Thr	Val	Asp	Cys	Asn	Asp	
				50					55						60
Leu	Arg	Leu	Thr	Arg	Ile	Pro	Ser	Asn	Leu	Ser	Ser	Asp	Thr	Gln	
				65					70						75
Val	Leu	Leu	Leu	Gln	Ser	Asn	Asn	Ile	Ala	Lys	Thr	Val	Asp	Glu	
				80					85						90
Leu	Gln	Gln	Leu	Phe	Asn	Leu	Thr	Glu	Leu	Asp	Phe	Ser	Gln	Asn	
				95					100						105
Asn	Phe	Thr	Asn	Ile	Lys	Glu	Val	Gly	Leu	Ala	Asn	Leu	Thr	Gln	
				110					115						120
Leu	Thr	Thr	Leu	His	Leu	Glu	Glu	Asn	Gln	Ile	Thr	Glu	Met	Thr	
				125					130						135
Asp	Tyr	Cys	Leu	Gln	Asp	Leu	Ser	Asn	Leu	Gln	Glu	Leu	Tyr	Ile	
				140					145						150
Asn	His	Asn	Gln	Ile	Ser	Thr	Ile	Ser	Ala	His	Ala	Phe	Ala	Gly	
				155					160						165
Leu	Lys	Asn	Leu	Leu	Arg	Leu	His	Leu	Asn	Ser	Asn	Lys	Leu	Lys	
				170					175						180
Val	Ile	Asp	Ser	Arg	Trp	Phe	Asp	Ser	Thr	Pro	Asn	Leu	Glu	Ile	
				185					190						195
Leu	Met	Ile	Gly	Glu	Asn	Pro	Val	Ile	Gly	Ile	Leu	Asp	Met	Asn	
				200					205						210
Phe	Lys	Pro	Leu	Ala	Asn	Leu	Arg	Ser	Leu	Val	Leu	Ala	Gly	Met	
				215					220						225
Tyr	Leu	Thr	Asp	Ile	Pro	Gly	Asn	Ala	Leu	Val	Gly	Leu	Asp	Ser	
				230					235						240
Leu	Glu	Ser	Leu	Ser	Phe	Tyr	Asp	Asn	Lys	Leu	Val	Lys	Val	Pro	
				245					250						255
Gln	Leu	Ala	Leu	Gln	Lys	Val	Pro	Asn	Leu	Lys	Phe	Leu	Asp	Leu	
				260					265						270
Asn	Lys	Asn	Pro	Ile	His	Lys	Ile	Gln	Glu	Gly	Asp	Phe	Lys	Asn	
				275					280						285
Met	Leu	Arg	Leu	Lys	Glu	Leu	Gly	Ile	Asn	Asn	Met	Gly	Glu	Leu	
				290					295						300
Val	Ser	Val	Asp	Arg	Tyr	Ala	Leu	Asp	Asn	Leu	Pro	Glu	Leu	Thr	
				305					310						315
Lys	Leu	Glu	Ala	Thr	Asn	Asn	Pro	Lys	Leu	Ser	Tyr	Ile	His	Arg	
				320					325						330
Leu	Ala	Phe	Arg	Ser	Val	Pro	Ala	Leu	Glu	Ser	Leu	Met	Leu	Asn	
				335					340						345
Asn	Asn	Ala	Leu	Asn	Ala	Ile	Tyr	Gln	Lys	Thr	Val	Glu	Ser	Leu	
				350					355						360
Pro	Asn	Leu	Arg	Glu	Ile	Ser	Ile	His	Ser	Asn	Pro	Leu	Arg	Cys	
				365					370						375
Asp	Cys	Val	Ile	His	Trp	Ile	Asn	Ser	Asn	Lys	Thr	Asn	Ile	Arg	
				380					385						390
Phe	Met	Glu	Pro	Leu	Ser	Met	Phe	Cys	Ala	Met	Pro	Pro	Glu	Tyr	
				395					400						405
Lys	Gly	His	Gln	Val	Lys	Glu	Val	Leu	Ile	Gln	Asp	Ser	Ser	Glu	
				410					415						420
Gln	Cys	Leu	Pro	Met	Ile	Ser	His	Asp	Ser	Phe	Pro	Asn	Arg	Leu	
				425					430						435
Asn	Val	Asp	Ile	Gly	Thr	Thr	Val	Phe	Leu	Asp	Cys	Arg	Ala	Met	
				440					445						450
Ala	Glu	Pro	Glu	Pro	Glu	Ile	Tyr	Trp	Val	Thr	Pro	Ile	Gly	Asn	
				455					460						465
Lys	Ile	Thr	Val	Glu	Thr	Leu	Ser	Asp	Lys	Tyr	Lys	Leu	Ser	Ser	
				470					475						480
Glu	Gly	Thr	Leu	Glu	Ile	Ser	Asn	Ile	Gln	Ile	Glu	Asp	Ser	Gly	
				485					490						495
Arg	Tyr	Thr	Cys	Val	Ala	Gln	Asn	Val	Gln	Gly	Ala	Asp	Thr	Arg	
				500					505						510
Val	Ala	Thr	Ile	Lys	Val	Asn	Gly	Thr	Leu	Leu	Asp	Gly	Thr	Gln	
				515					520						525
Val	Leu	Lys	Ile	Tyr	Val	Lys	Gln	Thr	Glu	Ser	His	Ser	Ile	Leu	

Val	Ser	Trp	Lys	530	Val	Asn	Ser	Asn	Val	535	Thr	Ser	Asn	Leu	540
				545	Met	Lys	Ile	Asp	Asn	550	Pro	His	Ile	Thr	555
Trp	Ser	Ser	Ala	560	Thr	Met	Lys	Ile	Asp	565	Tyr	Asn	Leu	Thr	570
Thr	Ala	Arg	Val	575	Pro	Val	Asp	Val	His	580	Glu	Tyr	Asn	Leu	585
Leu	Gln	Pro	Ser	590	Thr	Asp	Tyr	Glu	Val	595	Cys	Leu	Thr	Val	600
Ile	His	Gln	Gln	605	Thr	Gln	Lys	Ser	Cys	610	Val	Asn	Val	Thr	615
Asn	Ala	Ala	Phe	620	Ala	Val	Asp	Ile	Ser	625	Gln	Glu	Thr	Ser	630
Ala	Leu	Ala	Ala	635	Val	Met	Gly	Ser	Met	640	Phe	Ala	Val	Ile	645
Ala	Ser	Ile	Ala	650	Val	Tyr	Phe	Ala	Lys	655	Phe	Lys	Arg	Lys	660
Tyr	His	His	Ser	665	Leu	Lys	Lys	Tyr	Met	670	Gln	Lys	Thr	Ser	675
Pro	Leu	Asn	Glu	680	Leu	Tyr	Pro	Pro	Leu	685	Ile	Asn	Leu	Trp	690
Asp	Ser	Glu	Lys	695	Asp	Lys	Asp	Gly	Ser	700	Ala	Asp	Thr	Lys	705
Gln	Val	Asp	Thr	710	Ser	Arg	Ser	Tyr	Tyr	715	Met	Trp			

<210> 13

<211> 665

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4568384CD1

<400> 13

Met	Val	Leu	Val	Phe	His	Lys	Gly	Glu	Leu	Gly	His	Pro	Leu	Glu	
1				5					10					15	
Gln	Ser	Thr	Asp	Trp	Pro	Lys	Ser	Pro	Lys	Thr	Pro	Thr	Gly	Leu	
				20					25					30	
Arg	Arg	Gly	Arg	Gln	Cys	Ile	Arg	Pro	Ala	Glu	Ile	Val	Ala	Ser	
				35					40					45	
Leu	Leu	Glu	Gly	Glu	Glu	Asn	Thr	Cys	Gly	Lys	Gln	Lys	Pro	Lys	
				50					55					60	
Glu	Asn	Asn	Leu	Lys	Pro	Lys	Phe	Gln	Ala	Phe	Lys	Gly	Val	Gly	
				65					70					75	
Cys	Leu	Tyr	Glu	Lys	Glu	Ser	Met	Lys	Lys	Ser	Leu	Lys	Asp	Ser	
				80					85					90	
Val	Ala	Ser	Asn	Asn	Lys	Asp	Gln	Asn	Ser	Met	Lys	His	Glu	Asp	
				95					100					105	
Pro	Ser	Ile	Ile	Ser	Met	Glu	Asp	Gly	Ser	Pro	Tyr	Val	Asn	Gly	
				110					115					120	
Ser	Leu	Gly	Glu	Val	Thr	Pro	Cys	Gln	His	Ala	Lys	Lys	Ala	Asn	
				125					130					135	
Gly	Pro	Asn	Tyr	Ile	Gln	Pro	Gln	Lys	Arg	Gln	Thr	Thr	Phe	Glu	
				140					145					150	
Ser	Gln	Asp	Arg	Lys	Ala	Val	Ser	Pro	Ser	Ser	Ser	Glu	Lys	Arg	
				155					160					165	
Ser	Lys	Asn	Pro	Ile	Ser	Arg	Pro	Leu	Glu	Gly	Lys	Lys	Ser	Leu	
				170					175					180	
Ser	Leu	Ser	Ala	Lys	Thr	His	Asn	Ile	Gly	Phe	Asp	Lys	Asp	Ser	
				185					190					195	
Cys	His	Ser	Thr	Thr	Lys	Thr	Glu	Ala	Ser	Gln	Glu	Glu	Arg	Ser	
				200					205					210	
Asp	Ser	Ser	Gly	Leu	Thr	Ser	Leu	Lys	Lys	Ser	Pro	Lys	Val	Ser	
				215					220					225	
Ser	Lys	Asp	Thr	Arg	Glu	Ile	Lys	Thr	Asp	Phe	Ser	Leu	Ser	Ile	
				230					235					240	

Ser	Asn	Ser	Ser	Asp	Val	Ser	Ala	Lys	Asp	Lys	His	Ala	Glu	Asp
				245					250					255
Asn	Glu	Lys	Arg	Leu	Ala	Ala	Leu	Glu	Ala	Arg	Gln	Lys	Ala	Lys
				260					265					270
Glu	Val	Gln	Lys	Lys	Leu	Val	His	Asn	Ala	Leu	Ala	Asn	Leu	Asp
				275					280					285
Gly	His	Pro	Glu	Asp	Lys	Pro	Thr	His	Ile	Ile	Phe	Gly	Ser	Asp
				290					295					300
Ser	Glu	Cys	Glu	Thr	Glu	Glu	Thr	Ser	Thr	Gln	Glu	Gln	Ser	His
				305					310					315
Pro	Gly	Glu	Glu	Trp	Val	Lys	Glu	Ser	Met	Gly	Lys	Thr	Ser	Gly
				320					325					330
Lys	Leu	Phe	Asp	Ser	Ser	Asp	Asp	Asp	Glu	Ser	Asp	Ser	Glu	Asp
				335					340					345
Asp	Ser	Asn	Arg	Phe	Lys	Ile	Lys	Pro	Gln	Phe	Glu	Gly	Arg	Ala
				350					355					360
Gly	Gln	Lys	Leu	Met	Asp	Leu	Gln	Ser	His	Phe	Gly	Thr	Asp	Asp
				365					370					375
Arg	Phe	Arg	Met	Asp	Ser	Arg	Phe	Leu	Glu	Thr	Asp	Ser	Glu	Glu
				380					385					390
Glu	Gln	Glu	Glu	Val	Asn	Glu	Lys	Lys	Thr	Ala	Glu	Glu	Glu	Glu
				395					400					405
Leu	Ala	Glu	Glu	Lys	Lys	Lys	Ala	Leu	Asn	Val	Val	Gln	Ser	Val
				410					415					420
Leu	Gln	Ile	Asn	Leu	Ser	Asn	Ser	Thr	Asn	Arg	Gly	Ser	Val	Ala
				425					430					435
Ala	Lys	Lys	Phe	Lys	Asp	Ile	Ile	His	Tyr	Asp	Pro	Thr	Lys	Gln
				440					445					450
Asp	His	Ala	Thr	Tyr	Glu	Arg	Lys	Arg	Asp	Asp	Lys	Pro	Lys	Glu
				455					460					465
Ser	Lys	Ala	Lys	Arg	Lys	Lys	Lys	Arg	Glu	Glu	Ala	Glu	Lys	Leu
				470					475					480
Pro	Glu	Val	Ser	Lys	Glu	Met	Tyr	Tyr	Asn	Ile	Ala	Met	Asp	Leu
				485					490					495
Lys	Glu	Ile	Phe	Gln	Thr	Thr	Lys	Tyr	Thr	Ser	Glu	Lys	Glu	Glu
				500					505					510
Gly	Thr	Pro	Trp	Asn	Glu	Asp	Cys	Gly	Lys	Glu	Lys	Pro	Glu	Glu
				515					520					525
Ile	Gln	Asp	Pro	Ala	Ala	Leu	Thr	Ser	Asp	Ala	Glu	Gln	Pro	Ser
				530					535					540
Gly	Phe	Thr	Phe	Ser	Phe	Phe	Asp	Ser	Asp	Thr	Lys	Asp	Ile	Lys
				545					550					555
Glu	Glu	Thr	Tyr	Arg	Val	Glu	Thr	Val	Lys	Pro	Gly	Lys	Ile	Val
				560					565					570
Trp	Gln	Glu	Asp	Pro	Arg	Leu	Gln	Asp	Ser	Ser	Ser	Glu	Glu	Glu
				575					580					585
Asp	Val	Thr	Glu	Glu	Thr	Asp	His	Arg	Asn	Ser	Ser	Pro	Gly	Glu
				590					595					600
Ala	Ser	Leu	Leu	Glu	Lys	Glu	Thr	Thr	Arg	Phe	Phe	Phe	Phe	Ser
				605					610					615
Lys	Asn	Asp	Glu	Arg	Leu	Gln	Gly	Ser	Asp	Leu	Phe	Trp	Arg	Gly
				620					625					630
Val	Gly	Ser	Asn	Met	Ser	Arg	Asn	Ser	Trp	Glu	Ala	Arg	Thr	Thr
				635					640					645
Asn	Leu	Arg	Met	Asp	Cys	Arg	Lys	Lys	His	Lys	Asp	Ala	Lys	Arg
				650					655					660
Lys	Met	Lys	Pro	Lys										
				665										

<210> 14

<211> 547

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4586187CD1

<400> 14

Met	Tyr	Ser	His	Asn	Val	Val	Ile	Met	Asn	Leu	Asn	Asn	Leu	Asn
1				5					10					15
Leu	Thr	Gln	Val	Gln	Gln	Arg	Asn	Leu	Ile	Thr	Asn	Leu	Gln	Arg
				20					25					30
Ser	Val	Asp	Asp	Thr	Ser	Gln	Ala	Ile	Gln	Arg	Ile	Lys	Asn	Asp
				35					40					45
Phe	Gln	Asn	Leu	Gln	Gln	Val	Phe	Leu	Gln	Ala	Lys	Lys	Asp	Thr
				50					55					60
Asp	Trp	Leu	Lys	Glu	Lys	Val	Gln	Ser	Leu	Gln	Thr	Leu	Ala	Ala
				65					70					75
Asn	Asn	Ser	Ala	Leu	Ala	Lys	Ala	Asn	Asn	Asp	Thr	Leu	Glu	Asp
				80					85					90
Met	Asn	Ser	Gln	Leu	Asn	Ser	Phe	Thr	Gly	Gln	Met	Glu	Asn	Ile
				95					100					105
Thr	Thr	Ile	Ser	Gln	Ala	Asn	Glu	Gln	Asn	Leu	Lys	Asp	Leu	Gln
				110					115					120
Asp	Leu	His	Lys	Asp	Ala	Glu	Asn	Arg	Thr	Ala	Ile	Lys	Phe	Asn
				125					130					135
Gln	Leu	Glu	Glu	Arg	Phe	Gln	Leu	Phe	Glu	Thr	Asp	Ile	Val	Asn
				140					145					150
Ile	Ile	Ser	Asn	Ile	Ser	Tyr	Thr	Ala	His	His	Leu	Arg	Thr	Leu
				155					160					165
Thr	Ser	Asn	Leu	Asn	Glu	Val	Arg	Thr	Thr	Cys	Thr	Asp	Thr	Leu
				170					175					180
Thr	Lys	His	Thr	Asp	Asp	Leu	Thr	Ser	Leu	Asn	Asn	Thr	Leu	Ala
				185					190					195
Asn	Ile	Arg	Leu	Asp	Ser	Val	Ser	Leu	Arg	Met	Gln	Gln	Asp	Leu
				200					205					210
Met	Arg	Ser	Arg	Leu	Asp	Thr	Glu	Val	Ala	Asn	Leu	Ser	Val	Ile
				215					220					225
Met	Glu	Glu	Met	Lys	Leu	Val	Asp	Ser	Lys	His	Gly	Gln	Leu	Ile
				230					235					240
Lys	Asn	Phe	Thr	Ile	Leu	Gln	Gly	Pro	Pro	Gly	Pro	Arg	Gly	Pro
				245					250					255
Arg	Gly	Asp	Arg	Gly	Ser	Gln	Gly	Pro	Pro	Gly	Pro	Thr	Gly	Asn
				260					265					270
Lys	Gly	Gln	Lys	Gly	Glu	Lys	Gly	Glu	Pro	Gly	Pro	Pro	Gly	Pro
				275					280					285
Ala	Gly	Glu	Arg	Gly	Pro	Ile	Gly	Pro	Ala	Gly	Pro	Pro	Gly	Glu
				290					295					300
Arg	Gly	Gly	Lys	Gly	Ser	Lys	Gly	Ser	Gln	Gly	Pro	Lys	Gly	Ser
				305					310					315
Arg	Gly	Ser	Pro	Gly	Lys	Pro	Gly	Pro	Gln	Gly	Pro	Ser	Gly	Asp
				320					325					330
Pro	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Lys	Glu	Gly	Leu	Pro	Gly	Pro
				335					340					345
Gln	Gly	Pro	Pro	Gly	Phe	Gln	Gly	Leu	Gln	Gly	Thr	Val	Gly	Glu
				350					355					360
Pro	Gly	Val	Pro	Gly	Pro	Arg	Gly	Leu	Pro	Gly	Leu	Pro	Gly	Val
				365					370					375
Pro	Gly	Met	Pro	Gly	Pro	Lys	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro
				380					385					390
Ser	Gly	Ala	Val	Val	Pro	Leu	Ala	Leu	Gln	Asn	Glu	Pro	Thr	Pro
				395					400					405
Ala	Pro	Glu	Asp	Asn	Ser	Cys	Pro	Pro	His	Trp	Lys	Asn	Phe	Thr
				410					415					420
Asp	Lys	Cys	Tyr	Tyr	Phe	Ser	Val	Glu	Lys	Glu	Ile	Phe	Glu	Asp
				425					430					435
Ala	Lys	Leu	Phe	Cys	Glu	Asp	Lys	Ser	Ser	His	Leu	Val	Phe	Ile
				440					445					450
Asn	Thr	Arg	Glu	Glu	Gln	Gln	Trp	Ile	Lys	Lys	Gln	Met	Val	Gly
				455					460					465
Arg	Glu	Ser	His	Trp	Ile	Gly	Leu	Thr	Asp	Ser	Glu	Arg	Glu	Asn
				470					475					480
Glu	Trp	Lys	Trp	Leu	Asp	Gly	Thr	Ser	Pro	Asp	Tyr	Lys	Asn	Trp
				485					490					495

Lys	Ala	Gly	Gln	Pro	Asp	Asn	Trp	Gly	His	Gly	His	Gly	Pro	Gly
				500					505					510
Glu	Asp	Cys	Ala	Gly	Leu	Ile	Tyr	Ala	Gly	Gln	Trp	Asn	Asp	Phe
				515					520					525
Gln	Cys	Glu	Asp	Val	Asn	Asn	Phe	Ile	Cys	Glu	Lys	Asp	Arg	Glu
				530					535					540
Thr	Val	Leu	Ser	Ser	Ala	Leu								
				545										

<210> 15
 <211> 109
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 401801CD1

<400>	15													
Met	Tyr	Phe	Asn	Leu	Gln	Glu	Asn	Ile	Phe	Met	Tyr	Gly	Gly	Arg
1				5					10					15
Ile	Glu	Thr	Asn	Asp	Gly	Asn	Val	Thr	Asp	Glu	Leu	Trp	Val	Phe
				20					25					30
Asn	Ile	His	Ser	Gln	Ser	Trp	Ser	Thr	Lys	Thr	Pro	Thr	Val	Leu
				35					40					45
Gly	His	Gly	Gln	Gln	Tyr	Ala	Val	Glu	Gly	His	Ser	Ala	His	Ile
				50					55					60
Met	Glu	Leu	Asp	Ser	Arg	Asp	Val	Val	Met	Ile	Ile	Ile	Phe	Gly
				65					70					75
Tyr	Ser	Ala	Ile	Tyr	Gly	Tyr	Thr	Ser	Ser	Ile	Gln	Glu	Tyr	His
				80					85					90
Ile	Cys	Glu	Leu	Leu	Lys	Asn	Cys	Asn	Phe	Phe	Ile	Asp	Trp	Glu
				95					100					105
Cys	Phe	Ser	Leu											

<210> 16
 <211> 192
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1721842CD1

<400>	16													
Met	Asn	Lys	Arg	Asp	Tyr	Met	Asn	Thr	Ser	Val	Gln	Glu	Pro	Pro
1				5					10					15
Leu	Asp	Tyr	Ser	Phe	Arg	Ser	Ile	His	Val	Ile	Gln	Asp	Leu	Val
				20					25					30
Asn	Glu	Glu	Pro	Arg	Thr	Gly	Leu	Arg	Pro	Leu	Lys	Arg	Ser	Lys
				35					40					45
Ser	Gly	Lys	Ser	Leu	Thr	Gln	Ser	Leu	Trp	Leu	Asn	Asn	Asn	Val
				50					55					60
Leu	Asn	Asp	Leu	Arg	Asp	Phe	Asn	Gln	Val	Ala	Ser	Gln	Leu	Leu
				65					70					75
Glu	His	Pro	Glu	Asn	Leu	Ala	Trp	Ile	Asp	Leu	Ser	Phe	Asn	Asp
				80					85					90
Leu	Thr	Ser	Ile	Asp	Pro	Val	Leu	Thr	Thr	Phe	Phe	Asn	Leu	Ser
				95					100					105
Val	Leu	Tyr	Leu	His	Gly	Asn	Ser	Ile	Gln	Arg	Leu	Gly	Glu	Val
				110					115					120
Asn	Lys	Leu	Ala	Val	Leu	Pro	Arg	Leu	Arg	Ser	Leu	Thr	Leu	His
				125					130					135
Gly	Asn	Pro	Met	Glu	Glu	Glu	Lys	Gly	Tyr	Arg	Gln	Tyr	Val	Leu
				140					145					150
Cys	Thr	Leu	Ser	Arg	Ile	Thr	Thr	Phe	Asp	Phe	Ser	Gly	Val	Thr
				155					160					165
Lys	Ala	Asp	Arg	Thr	Thr	Ala	Glu	Val	Trp	Lys	Arg	Met	Asn	Ile

```

170
Lys Pro Lys Lys Ala Trp Thr Lys Gln Asn Thr Leu
185
175
190
180
<210> 17
<211> 575
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1833221CD1

<400> 17
Met Val Leu Gly Ser Phe Gly Thr Asp Leu Met Arg Glu Arg Arg
1 5 10 15
Asp Leu Glu Arg Arg Thr Asp Ser Ser Ile Ser Asn Leu Met Asp
20 25 30
Tyr Ser His Arg Ser Gly Asp Phe Thr Thr Ser Ser Tyr Val Gln
35 40 45
Asp Arg Val Pro Ser Tyr Ser Gln Gly Ala Arg Pro Lys Glu Asn
50 55 60
Ser Met Ser Thr Leu Gln Leu Asn Thr Ser Ser Thr Asn His Gln
65 70 75
Leu Pro Ser Glu His Gln Thr Ile Leu Ser Ser Arg Asp Ser Arg
80 85 90
Asn Ser Leu Arg Ser Asn Phe Ser Ser Arg Glu Ser Glu Ser Ser
95 100 105
Arg Ser Asn Thr Gln Pro Gly Phe Ser Tyr Ser Ser Ser Arg Asp
110 115 120
Glu Ala Pro Ile Ile Ser Asn Ser Glu Arg Val Val Ser Ser Gln
125 130 135
Arg Pro Phe Gln Glu Ser Ser Asp Asn Glu Gly Arg Arg Thr Thr
140 145 150
Arg Arg Leu Leu Ser Arg Ile Ala Ser Ser Met Ser Ser Thr Phe
155 160 165
Phe Ser Arg Arg Ser Ser Gln Asp Ser Leu Asn Thr Arg Ser Leu
170 175 180
Asn Ser Glu Asn Ser Tyr Val Ser Pro Arg Ile Leu Thr Ala Ser
185 190 195
Gln Ser Arg Ser Asn Val Pro Ser Ala Ser Glu Val Pro Asp Asn
200 205 210
Arg Ala Ser Glu Ala Ser Gln Gly Phe Arg Phe Leu Arg Arg Arg
215 220 225
Trp Gly Leu Ser Ser Leu Ser His Asn His Ser Ser Glu Ser Asp
230 235 240
Ser Glu Asn Phe Asn Gln Glu Ser Glu Gly Arg Asn Thr Gly Pro
245 250 255
Trp Leu Ser Ser Ser Leu Arg Asn Arg Cys Thr Pro Leu Phe Ser
260 265 270
Arg Arg Arg Arg Glu Gly Arg Asp Glu Ser Ser Arg Ile Pro Thr
275 280 285
Ser Asp Thr Ser Ser Arg Ser His Ile Phe Arg Arg Glu Ser Asn
290 295 300
Glu Val Val His Leu Glu Ala Gln Asn Asp Pro Leu Gly Ala Ala
305 310 315
Ala Asn Arg Pro Gln Ala Ser Ala Ala Ser Ser Ser Ala Thr Thr
320 325 330
Gly Gly Ser Thr Ser Asp Ser Ala Gln Gly Gly Arg Asn Thr Gly
335 340 345
Ile Ser Gly Ile Leu Pro Gly Ser Leu Phe Arg Phe Ala Val Pro
350 355 360
Pro Ala Leu Gly Ser Asn Leu Thr Asp Asn Val Met Ile Thr Val
365 370 375
Asp Ile Ile Pro Ser Gly Trp Asn Ser Ala Asp Gly Lys Ser Asp
380 385 390
Lys Thr Lys Ser Ala Pro Ser Arg Asp Pro Glu Arg Leu Gln Lys
395 400 405

```

Ile	Lys	Glu	Ser	Leu	Leu	Glu	Asp	Ser	Glu	Glu	Glu	Glu	Gly
				410				415					420
Asp	Leu	Cys	Arg	Ile	Cys	Gln	Met	Ala	Ala	Ala	Ser	Ser	Asn
				425				430					435
Leu	Leu	Ile	Glu	Pro	Cys	Lys	Cys	Thr	Gly	Ser	Leu	Gln	Val
				440				445					450
His	Gln	Asp	Cys	Met	Lys	Lys	Trp	Leu	Gln	Ala	Lys	Ile	Ser
				455				460					465
Gly	Ser	Ser	Leu	Glu	Ala	Val	Thr	Thr	Cys	Glu	Leu	Cys	Glu
				470				475					480
Lys	Leu	Glu	Leu	Asn	Leu	Glu	Asp	Phe	Asp	Ile	His	Glu	His
				485				490					495
Arg	Ala	His	Ala	Asn	Glu	Gln	Ala	Glu	Tyr	Glu	Phe	Ile	Ser
				500				505					510
Gly	Leu	Tyr	Leu	Val	Val	Leu	Leu	His	Leu	Cys	Glu	Gln	Phe
				515				520					525
Ser	Asp	Met	Met	Gly	Asn	Thr	Asn	Glu	Pro	Ser	Thr	Arg	Val
				530				535					540
Phe	Ile	Asn	Leu	Ala	Arg	Thr	Leu	Gln	Ala	His	Met	Glu	Asp
				545				550					555
Glu	Thr	Ser	Glu	Asp	Asp	Ser	Glu	Glu	Asp	Gly	Asp	His	Asn
				560				565					570
Thr	Phe	Asp	Ile	Ala									
				575									

<210> 18

<211> 342

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2041168CD1

<400> 18

Met	Ala	Glu	Gly	Gly	Ser	Gly	Asp	Val	Asp	Asp	Ala	Gly	Asp	Cys
1				5					10					15
Ser	Gly	Ala	Arg	Tyr	Asn	Asp	Trp	Ser	Asp	Asp	Asp	Asp	Asp	Ser
				20					25					30
Asn	Glu	Ser	Lys	Ser	Ile	Val	Trp	Tyr	Pro	Pro	Trp	Ala	Arg	Ile
				35					40					45
Gly	Thr	Glu	Ala	Gly	Thr	Arg	Ala	Arg	Ala	Arg	Ala	Arg	Ala	Arg
				50					55					60
Ala	Thr	Arg	Ala	Arg	Arg	Ala	Val	Gln	Lys	Arg	Ala	Ser	Pro	Asn
				65					70					75
Ser	Asp	Asp	Thr	Val	Leu	Ser	Pro	Gln	Glu	Leu	Gln	Lys	Val	Leu
				80					85					90
Cys	Leu	Val	Glu	Met	Ser	Glu	Lys	Pro	Tyr	Ile	Leu	Glu	Ala	Ala
				95					100					105
Leu	Ile	Ala	Leu	Gly	Asn	Asn	Ala	Ala	Tyr	Ala	Phe	Asn	Arg	Asp
				110					115					120
Ile	Ile	Arg	Asp	Leu	Gly	Gly	Leu	Pro	Ile	Val	Ala	Lys	Ile	Leu
				125					130					135
Asn	Thr	Arg	Asp	Pro	Ile	Val	Lys	Glu	Lys	Ala	Leu	Ile	Val	Leu
				140					145					150
Asn	Asn	Leu	Ser	Val	Asn	Ala	Glu	Asn	Gln	Arg	Arg	Leu	Lys	Val
				155					160					165
Tyr	Met	Asn	Gln	Val	Cys	Asp	Asp	Thr	Ile	Thr	Ser	Arg	Leu	Asn
				170					175					180
Ser	Ser	Val	Gln	Leu	Ala	Gly	Leu	Arg	Leu	Leu	Thr	Asn	Met	Thr
				185					190					195
Val	Thr	Asn	Glu	Tyr	Gln	His	Met	Leu	Ala	Asn	Ser	Ile	Ser	Asp
				200					205					210
Phe	Phe	Arg	Leu	Phe	Ser	Ala	Gly	Asn	Glu	Glu	Thr	Lys	Leu	Gln
				215					220					225
Val	Leu	Lys	Leu	Leu	Leu	Asn	Leu	Ala	Glu	Asn	Pro	Ala	Met	Thr
				230					235					240
Arg	Glu	Leu	Leu	Arg	Ala	Gln	Val	Pro	Ser	Ser	Leu	Gly	Ser	Leu

Phe	Asn	Lys	Lys	245	Asn	Lys	Glu	Val	250	Ile	Leu	Lys	Leu	Leu	Val	255
				260					265							270
Ile	Phe	Glu	Asn	275	Ile	Asn	Asp	Asn	280	Phe	Lys	Trp	Glu	Glu	Asn	285
Pro	Thr	Gln	Asn	290	Gln	Phe	Gly	Glu	295	Gly	Ser	Leu	Phe	Phe	Phe	300
Lys	Glu	Phe	Gln	305	Val	Cys	Ala	Asp	310	Lys	Val	Leu	Gly	Ile	Glu	315
His	His	Asp	Phe	320	Leu	Val	Lys	Val	325	Lys	Val	Gly	Lys	Phe	Met	330
Lys	Leu	Ala	Glu	335	His	Met	Phe	Pro	340	Lys	Ser	Gln	Glu			

<210> 19
 <211> 110
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2365794CD1

Met	Ala	Ala	Val	Val	Ala	Lys	Arg	Glu	Gly	Pro	Pro	Phe	Ile	Ser	
1				5					10					15	
Glu	Ala	Ala	Val	Arg	Gly	Asn	Ala	Ala	Val	Leu	Asp	Tyr	Cys	Arg	
				20					25					30	
Thr	Ser	Val	Ser	Ala	Leu	Ser	Gly	Ala	Thr	Ala	Gly	Ile	Leu	Gly	
				35					40					45	
Leu	Thr	Gly	Leu	Tyr	Gly	Phe	Ile	Phe	Tyr	Leu	Leu	Ala	Ser	Val	
				50					55					60	
Leu	Leu	Ser	Leu	Leu	Leu	Ile	Leu	Lys	Ala	Gly	Arg	Arg	Trp	Asn	
				65					70					75	
Lys	Tyr	Phe	Lys	Ser	Arg	Arg	Pro	Leu	Phe	Thr	Gly	Gly	Leu	Ile	
				80					85					90	
Gly	Gly	Leu	Phe	Thr	Tyr	Val	Leu	Phe	Trp	Thr	Phe	Leu	Tyr	Gly	
				95					100					105	
Met	Val	His	Val	Tyr											
				110											

<210> 20
 <211> 571
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2618452CD1

Met	Pro	Thr	Gly	Thr	Ile	Pro	Pro	Pro	Thr	Thr	Leu	Lys	Ala	Thr	
1				5					10					15	
Gly	Ser	Thr	His	Thr	Ala	Pro	Pro	Met	Met	Pro	Thr	Thr	Ser	Gly	
				20					25					30	
Thr	Ser	Gln	Ala	Ser	Ser	Ser	Phe	Asn	Thr	Ala	Lys	Thr	Ser	Thr	
				35					40					45	
Ser	Leu	His	Ser	His	Thr	Ser	Ser	Thr	His	His	Pro	Glu	Val	Thr	
				50					55					60	
Pro	Thr	Ser	Ile	Thr	Asn	Ile	Thr	Leu	Asn	Pro	Thr	Ser	Ile	Gly	
				65					70					75	
Thr	Trp	Thr	Pro	Val	Ala	His	Thr	Thr	Ser	Ala	Thr	Ser	Ser	Arg	
				80					85					90	
Leu	Thr	Thr	Pro	Phe	Thr	Thr	His	Ser	Pro	Pro	Thr	Gly	Ser	Ser	
				95					100					105	
Pro	Ile	Ser	Ser	Thr	Gly	Pro	Met	Thr	Ala	Thr	Ser	Phe	Gln	Thr	
				110					115					120	
Thr	Thr	Tyr	Tyr	Thr	Pro	Pro	Ser	His	Pro	Gln	Thr	Thr	Leu	Pro	
				125					130					135	

Thr	His	Val	Pro	Pro	Phe	Ser	Thr	Ser	Leu	Val	Thr	Pro	Ser	Thr	140	145	150
His	Thr	Val	Ile	Ile	Thr	Thr	His	Thr	Gln	Met	Ala	Thr	Ser	Ala	155	160	165
Ser	Ile	His	Ser	Thr	Pro	Thr	Gly	Thr	Val	Pro	Pro	Pro	Thr	Thr	170	175	180
Leu	Lys	Ala	Thr	Gly	Ser	Thr	His	Thr	Ala	Pro	Pro	Met	Thr	Val	185	190	195
Thr	Thr	Ser	Gly	Thr	Ser	Gln	Thr	His	Ser	Ser	Phe	Ser	Thr	Ala	200	205	210
Thr	Ala	Ser	Ser	Ser	Phe	Ile	Ser	Ser	Ser	Ser	Trp	Ser	Ser	Trp	215	220	225
Leu	Pro	Gln	Asn	Ser	Ser	Ser	Arg	Pro	Pro	Ser	Ser	Pro	Ile	Thr	230	235	240
Thr	Gln	Leu	Pro	His	Leu	Ser	Ser	Ala	Thr	Thr	Pro	Val	Ser	Thr	245	250	255
Thr	Asn	Gln	Leu	Ser	Ser	Ser	Phe	Ser	Pro	Ser	Pro	Ser	Ala	Pro	260	265	270
Ser	Thr	Val	Ser	Ser	Tyr	Val	Pro	Ser	Ser	His	Ser	Ser	Pro	Gln	275	280	285
Thr	Ser	Ser	Pro	Ser	Val	Gly	Thr	Ser	Ser	Ser	Phe	Val	Ser	Ala	290	295	300
Pro	Val	His	Ser	Thr	Thr	Leu	Ser	Ser	Gly	Ser	His	Ser	Ser	Leu	305	310	315
Ser	Thr	His	Pro	Thr	Thr	Ala	Ser	Val	Ser	Ala	Ser	Pro	Leu	Phe	320	325	330
Pro	Ser	Ser	Pro	Ala	Ala	Ser	Thr	Thr	Ile	Arg	Ala	Thr	Leu	Pro	335	340	345
His	Thr	Ile	Ser	Ser	Pro	Phe	Thr	Leu	Ser	Ala	Leu	Leu	Pro	Ile	350	355	360
Ser	Thr	Val	Thr	Val	Ser	Pro	Thr	Pro	Ser	Ser	His	Leu	Ala	Ser	365	370	375
Ser	Thr	Ile	Ala	Phe	Pro	Ser	Thr	Pro	Arg	Thr	Thr	Ala	Ser	Thr	380	385	390
His	Thr	Ala	Pro	Ala	Phe	Ser	Ser	Gln	Ser	Thr	Thr	Ser	Arg	Ser	395	400	405
Thr	Ser	Leu	Thr	Thr	Arg	Val	Pro	Thr	Ser	Gly	Phe	Val	Ser	Leu	410	415	420
Thr	Ser	Gly	Val	Thr	Gly	Ile	Pro	Thr	Ser	Pro	Val	Thr	Asn	Leu	425	430	435
Thr	Thr	Arg	His	Pro	Gly	Pro	Thr	Leu	Ser	Pro	Thr	Thr	Arg	Phe	440	445	450
Leu	Thr	Ser	Ser	Leu	Thr	Ala	His	Gly	Ser	Thr	Pro	Ala	Ser	Ala	455	460	465
Pro	Val	Ser	Ser	Leu	Gly	Thr	Pro	Thr	Pro	Thr	Ser	Pro	Gly	Val	470	475	480
Cys	Ser	Val	Arg	Glu	Gln	Gln	Glu	Glu	Ile	Thr	Phe	Lys	Gly	Cys	485	490	495
Met	Ala	Asn	Val	Thr	Val	Thr	Arg	Cys	Glu	Gly	Ala	Cys	Ile	Ser	500	505	510
Ala	Ala	Ser	Phe	Asn	Ile	Ile	Thr	Gln	Gln	Val	Asp	Ala	Arg	Cys	515	520	525
Ser	Cys	Cys	Arg	Pro	Leu	His	Ser	Tyr	Glu	Gln	Gln	Leu	Glu	Leu	530	535	540
Pro	Cys	Pro	Asp	Pro	Ser	Thr	Pro	Gly	Arg	Arg	Leu	Val	Leu	Thr	545	550	555
Leu	Gln	Val	Phe	Ser	His	Cys	Val	Cys	Ser	Ser	Val	Ala	Cys	Gly	560	565	570

Asp

<210> 21

<211> 262

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2622288CD1

<400> 21
Met Val Ala Trp Arg Ser Ala Phe Leu Val Cys Leu Ala Phe Ser
1 5 10 15
Leu Ala Thr Leu Val Gln Arg Gly Ser Gly Asp Phe Asp Asp Phe
20 25 30
Asn Leu Glu Asp Ala Val Lys Glu Thr Ser Ser Val Lys Gln Pro
35 40 45
Trp Asp His Thr Thr Thr Thr Thr Thr Asn Arg Pro Gly Thr Thr
50 55 60
Arg Ala Pro Ala Lys Pro Pro Gly Ser Gly Leu Asp Leu Ala Asp
65 70 75
Ala Leu Asp Asp Gln Asp Asp Gly Arg Arg Lys Pro Gly Ile Gly
80 85 90
Gly Arg Glu Arg Trp Asn His Val Thr Thr Thr Thr Lys Arg Pro
95 100 105
Val Thr Thr Arg Ala Pro Ala Asn Thr Leu Gly Asn Asp Phe Asp
110 115 120
Leu Ala Asp Ala Leu Asp Asp Arg Asn Asp Arg Asp Asp Gly Arg
125 130 135
Arg Lys Pro Ile Ala Gly Gly Gly Gly Phe Ser Asp Lys Asp Leu
140 145 150
Glu Asp Ile Val Gly Gly Gly Glu Tyr Lys Pro Asp Lys Gly Lys
155 160 165
Gly Asp Gly Arg Tyr Gly Ser Asn Asp Asp Pro Gly Ser Gly Met
170 175 180
Val Ala Glu Pro Gly Thr Ile Ala Gly Val Ala Ser Ala Leu Ala
185 190 195
Met Ala Leu Ile Gly Ala Val Ser Ser Tyr Ile Ser Tyr Gln Gln
200 205 210
Lys Lys Phe Cys Phe Ser Ile Gln Gln Gly Leu Asn Ala Asp Tyr
215 220 225
Val Lys Gly Glu Asn Leu Glu Ala Val Val Cys Glu Glu Pro Gln
230 235 240
Val Lys Tyr Ser Thr Leu His Thr Gln Ser Ala Glu Pro Pro Pro
245 250 255
Pro Pro Glu Pro Ala Arg Ile
260

<210> 22
<211> 172
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2806595CD1

<400> 22
Met Gly Leu Leu Leu Leu Val Pro Leu Leu Leu Leu Pro Gly Ser
1 5 10 15
Tyr Gly Leu Pro Phe Tyr Asn Gly Phe Tyr Tyr Ser Asn Ser Ala
20 25 30
Asn Asp Gln Asn Leu Gly Asn Gly His Gly Lys Asp Leu Leu Asn
35 40 45
Gly Val Lys Leu Val Val Glu Thr Pro Glu Glu Thr Leu Phe Thr
50 55 60
Tyr Gln Gly Ala Ser Val Ile Leu Pro Cys Arg Tyr Arg Tyr Glu
65 70 75
Pro Ala Leu Val Ser Pro Arg Arg Val Arg Val Lys Trp Trp Lys
80 85 90
Leu Ser Glu Asn Gly Ala Pro Glu Lys Asp Val Leu Val Ala Ile
95 100 105
Gly Leu Arg His Arg Ser Phe Gly Asp Tyr Gln Gly Arg Val His
110 115 120
Leu Arg Gln Asp Lys Glu His Asp Val Ser Leu Glu Ile Gln Asp
125 130 135
Leu Arg Leu Glu Asp Tyr Gly Arg Tyr Arg Cys Glu Val Ile Asp
140 145 150

Gly Leu Glu Asp Glu Ser Gly Leu Val Glu Leu Glu Leu Arg Gly
 155 160 165
 Glu Met Leu Thr Gly Thr Gly
 170

<210> 23

<211> 571

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2850987CD1

<400> 23

Met Thr Arg Ala Gly Asp His Asn Arg Gln Arg Gly Cys Cys Gly
 1 5 10 15
 Ser Leu Ala Asp Tyr Leu Thr Ser Ala Lys Phe Leu Leu Tyr Leu
 20 25 30
 Gly His Ser Leu Ser Thr Trp Gly Asp Arg Met Trp His Phe Ala
 35 40 45
 Val Ser Val Phe Leu Val Glu Leu Tyr Gly Asn Ser Leu Leu Leu
 50 55 60
 Thr Ala Val Tyr Gly Leu Val Val Ala Gly Ser Val Leu Val Leu
 65 70 75
 Gly Ala Ile Ile Gly Asp Trp Val Asp Lys Asn Ala Arg Leu Lys
 80 85 90
 Val Ala Gln Thr Ser Leu Val Val Gln Asn Val Ser Val Ile Leu
 95 100 105
 Cys Gly Ile Ile Leu Met Met Val Phe Leu His Lys His Glu Leu
 110 115 120
 Leu Thr Met Tyr His Gly Trp Val Leu Thr Ser Cys Tyr Ile Leu
 125 130 135
 Ile Ile Thr Ile Ala Asn Ile Ala Asn Leu Ala Ser Thr Ala Thr
 140 145 150
 Ala Ile Thr Ile Gln Arg Asp Trp Ile Val Val Val Ala Gly Glu
 155 160 165
 Asp Arg Ser Lys Leu Ala Asn Met Asn Ala Thr Ile Arg Arg Ile
 170 175 180
 Asp Gln Leu Thr Asn Ile Leu Ala Pro Met Ala Val Gly Gln Ile
 185 190 195
 Met Thr Phe Gly Ser Pro Val Ile Gly Cys Gly Phe Ile Ser Gly
 200 205 210
 Trp Asn Leu Val Ser Met Cys Val Glu Tyr Val Leu Leu Trp Lys
 215 220 225
 Val Tyr Gln Lys Thr Pro Ala Leu Ala Val Lys Ala Gly Leu Lys
 230 235 240
 Glu Glu Glu Thr Glu Leu Lys Gln Leu Asn Leu His Lys Asp Thr
 245 250 255
 Glu Pro Lys Pro Leu Glu Gly Thr His Leu Met Gly Val Lys Asp
 260 265 270
 Ser Asn Ile His Glu Leu Glu His Glu Gln Glu Pro Thr Cys Ala
 275 280 285
 Ser Gln Met Ala Glu Pro Phe Arg Thr Phe Arg Asp Gly Trp Val
 290 295 300
 Ser Tyr Tyr Asn Gln Pro Val Phe Leu Ala Gly Met Gly Leu Ala
 305 310 315
 Phe Leu Tyr Met Thr Val Leu Gly Phe Asp Cys Ile Thr Thr Gly
 320 325 330
 Tyr Ala Tyr Thr Gln Gly Leu Ser Gly Ser Ile Leu Ser Ile Leu
 335 340 345
 Met Gly Ala Ser Ala Ile Thr Gly Ile Met Gly Thr Val Ala Phe
 350 355 360
 Thr Trp Leu Arg Arg Lys Cys Gly Leu Val Arg Thr Gly Leu Ile
 365 370 375
 Ser Gly Leu Ala Gln Leu Ser Cys Leu Ile Leu Cys Val Ile Ser
 380 385 390
 Val Phe Met Pro Gly Ser Pro Leu Asp Leu Ser Val Ser Pro Phe

	395		400		405
Glu Asp Ile Arg	Ser Arg Phe Ile Gln	Gly Glu Ser Ile Thr	Pro		
	410		415		420
Thr Lys Ile Pro	Glu Ile Thr Thr Glu	Ile Tyr Met Ser Asn	Gly		
	425		430		435
Ser Asn Ser Ala	Asn Ile Val Pro Glu	Thr Ser Pro Glu Ser	Val		
	440		445		450
Pro Ile Ile Ser	Val Ser Leu Leu Phe	Ala Gly Val Ile Ala	Ala		
	455		460		465
Arg Ile Gly Leu	Trp Ser Phe Asp Leu	Thr Val Thr Gln Leu	Leu		
	470		475		480
Gln Glu Asn Val	Ile Glu Ser Glu Arg	Gly Ile Ile Asn Gly	Val		
	485		490		495
Gln Asn Ser Met	Asn Tyr Leu Leu Asp	Leu Leu His Phe Ile	Met		
	500		505		510
Val Ile Leu Ala	Pro Asn Pro Glu Ala	Phe Gly Leu Leu Val	Leu		
	515		520		525
Ile Ser Val Ser	Phe Val Ala Met Gly	His Ile Met Tyr Phe	Arg		
	530		535		540
Phe Ala Gln Asn	Thr Leu Gly Asn Lys	Leu Phe Ala Cys Gly	Pro		
	545		550		555
Asp Ala Lys Glu	Val Arg Lys Glu Asn	Gln Ala Asn Thr Ser	Val		
	560		565		570

Val

<210> 24

<211> 455

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3557211CD1

<400> 24

Met Asp Pro Thr	Gly Asn Ser Ala Thr	Pro Gln Ile Leu Glu	Leu
1	5	10	15
Lys Trp Ser His	Ile Glu Trp Ser Gln	Thr Glu Tyr Ile Cys	Glu
	20	25	30
Asn Val Gly Leu	Leu Pro Leu Glu Ile	Ile Arg Arg Gly Tyr	Ser
	35	40	45
Met Asp Ser Ala	Phe Val Gly Ile Lys	Val Asn Gln Val Ser	Ala
	50	55	60
Ala Val Gly Lys	Asp Phe Thr Val Ile	Pro Ser Lys Leu Ile	Gln
	65	70	75
Phe Asp Pro Gly	Met Ser Thr Lys Met	Trp Asn Ile Ala Ile	Thr
	80	85	90
Tyr Asp Gly Leu	Glu Glu Asp Asp Glu	Val Phe Glu Val Ile	Leu
	95	100	105
Asn Ser Pro Val	Asn Ala Val Leu Gly	Thr Lys Thr Lys Ala	Ala
	110	115	120
Val Lys Ile Leu	Asp Ser Lys Gly Gly	Gln Cys His Pro Ser	Tyr
	125	130	135
Ser Ser Asn Gln	Ser Lys His Ser Thr	Trp Glu Lys Gly Ile	Trp
	140	145	150
His Leu Leu Pro	Pro Gly Ser Ser Ser	Ser Thr Thr Ser Gly	Ser
	155	160	165
Phe His Leu Glu	Arg Arg Pro Leu Pro	Ser Ser Met Gln Leu	Ala
	170	175	180
Val Ile Arg Gly	Asp Thr Leu Arg Gly	Phe Asp Ser Thr Asp	Leu
	185	190	195
Ser Gln Arg Lys	Leu Arg Thr Arg Gly	Asn Gly Lys Thr Val	Arg
	200	205	210
Pro Ser Ser Val	Tyr Arg Asn Gly Thr	Asp Ile Ile Tyr Asn	Tyr
	215	220	225
His Gly Ile Val	Ser Leu Lys Leu Glu	Asp Asp Ser Phe Pro	Thr
	230	235	240
His Lys Arg Lys	Ala Lys Val Ser Ile	Ile Ser Gln Pro Gln	Lys

Thr	Ile	Lys	Val	245	Ala	Glu	Leu	Pro	Gln	250	Ala	Asp	Lys	Val	Glu	255
				260						265						270
Thr	Thr	Asp	Ser	275	His	Phe	Pro	Arg	Gln	280	Asp	Gln	Leu	Pro	Ser	285
Pro	Lys	Asn	Cys	290	Thr	Leu	Glu	Leu	Lys	295	Gly	Leu	Phe	His	Phe	300
Glu	Gly	Ile	Gln	305	Lys	Leu	Tyr	Gln	Cys	310	Asn	Gly	Ile	Ala	Trp	315
Ala	Trp	Ser	Pro	320	Gln	Thr	Lys	Asp	Val	325	Glu	Asp	Lys	Ser	Cys	330
Ala	Gly	Trp	His	335	Gln	His	Ser	Gly	Tyr	340	Cys	His	Ile	Leu	Ile	345
Glu	Gln	Lys	Gly	350	Thr	Trp	Asn	Ala	Ala	355	Ala	Gln	Ala	Cys	Arg	360
Gln	Tyr	Leu	Gly	365	Asn	Leu	Val	Thr	Val	370	Phe	Ser	Arg	Gln	His	375
Arg	Trp	Leu	Trp	380	Asp	Ile	Gly	Gly	Arg	385	Lys	Ser	Phe	Trp	Ile	390
Leu	Asn	Asp	Gln	395	Val	His	Ala	Gly	His	400	Trp	Glu	Trp	Ile	Gly	405
Glu	Pro	Val	Ala	410	Phe	Thr	Asn	Gly	Arg	415	Arg	Gly	Pro	Ser	Pro	420
Ser	Lys	Leu	Gly	425	Lys	Ser	Cys	Val	Leu	430	Val	Gln	Arg	Gln	Gly	435
Trp	Gln	Thr	Lys	440	Asp	Cys	Arg	Arg	Ala	445	Lys	Pro	His	Asn	Tyr	450
Cys	Ser	Arg	Lys	455	Leu											

<210> 25

<211> 437

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4675668CD1

<400> 25

Met	Pro	Lys	Phe	Lys	Ala	Ala	Arg	Gly	Val	Gly	Gly	Gln	Glu	Lys	
1				5					10					15	
His	Ala	Pro	Leu	Ala	Asp	Gln	Ile	Leu	Ala	Gly	Asn	Ala	Val	Arg	
				20					25					30	
Ala	Gly	Val	Arg	Glu	Lys	Arg	Arg	Gly	Arg	Gly	Thr	Gly	Glu	Ala	
				35					40					45	
Glu	Glu	Glu	Tyr	Val	Gly	Pro	Arg	Leu	Ser	Arg	Arg	Ile	Leu	Gln	
				50					55					60	
Gln	Ala	Arg	Gln	Gln	Gln	Glu	Glu	Leu	Glu	Ala	Glu	His	Gly	Thr	
				65					70					75	
Gly	Asp	Lys	Pro	Ala	Ala	Pro	Arg	Glu	Arg	Thr	Thr	Arg	Leu	Gly	
				80					85					90	
Pro	Arg	Met	Pro	Gln	Asp	Gly	Ser	Asp	Asp	Glu	Asp	Glu	Glu	Trp	
				95					100					105	
Pro	Thr	Leu	Glu	Lys	Ala	Ala	Thr	Met	Thr	Ala	Ala	Gly	His	His	
				110					115					120	
Ala	Glu	Val	Val	Val	Asp	Pro	Glu	Asp	Arg	Arg	Ala	Ile	Glu	Met	
				125					130					135	
Phe	Met	Asn	Lys	Asn	Pro	Pro	Ala	Arg	Arg	Thr	Leu	Ala	Asp	Ile	
				140					145					150	
Ile	Met	Glu	Lys	Leu	Thr	Glu	Lys	Gln	Thr	Glu	Val	Glu	Thr	Val	
				155					160					165	
Met	Ser	Glu	Val	Ser	Gly	Phe	Pro	Met	Pro	Gln	Leu	Asp	Pro	Arg	
				170					175					180	
Val	Leu	Glu	Val	Tyr	Arg	Gly	Val	Arg	Glu	Val	Leu	Ser	Lys	Tyr	
				185					190					195	
Arg	Ser	Gly	Lys	Leu	Pro	Lys	Ala	Phe	Lys	Ile	Ile	Pro	Ala	Leu	
				200					205					210	

Ser	Asn	Trp	Glu	Gln	Ile	Leu	Tyr	Val	Thr	Glu	Pro	Glu	Ala	Trp
				215					220					225
Thr	Ala	Ala	Ala	Met	Tyr	Gln	Ala	Thr	Arg	Ile	Phe	Ala	Ser	Asn
				230					235					240
Leu	Lys	Glu	Arg	Met	Ala	Gln	Arg	Phe	Tyr	Asn	Leu	Val	Leu	Leu
				245					250					255
Pro	Arg	Val	Arg	Asp	Asp	Val	Ala	Glu	Tyr	Lys	Arg	Leu	Asn	Phe
				260					265					270
His	Leu	Tyr	Met	Ala	Leu	Lys	Lys	Ala	Leu	Phe	Lys	Pro	Gly	Ala
				275					280					285
Trp	Phe	Lys	Gly	Ile	Leu	Ile	Pro	Leu	Cys	Glu	Ser	Gly	Thr	Cys
				290					295					300
Thr	Leu	Arg	Glu	Ala	Ile	Ile	Val	Gly	Ser	Ile	Ile	Thr	Lys	Cys
				305					310					315
Ser	Ile	Pro	Val	Leu	His	Ser	Ser	Ala	Ala	Met	Leu	Lys	Ile	Ala
				320					325					330
Glu	Met	Glu	Tyr	Ser	Gly	Ala	Asn	Ser	Ile	Phe	Leu	Arg	Leu	Leu
				335					340					345
Leu	Asp	Lys	Lys	Tyr	Ala	Leu	Pro	Tyr	Arg	Val	Leu	Asp	Ala	Leu
				350					355					360
Val	Phe	His	Phe	Leu	Gly	Phe	Arg	Thr	Glu	Lys	Arg	Glu	Leu	Pro
				365					370					375
Val	Leu	Trp	His	Gln	Cys	Leu	Leu	Thr	Leu	Val	Gln	Arg	Tyr	Lys
				380					385					390
Ala	Asp	Leu	Ala	Thr	Asp	Gln	Lys	Glu	Ala	Leu	Leu	Glu	Leu	Leu
				395					400					405
Arg	Leu	Gln	Pro	His	Pro	Gln	Leu	Ser	Pro	Glu	Ile	Arg	Arg	Glu
				410					415					420
Leu	Gln	Ser	Ala	Val	Pro	Arg	Asp	Val	Glu	Asp	Val	Pro	Ile	Thr
				425					430					435

Val Glu

<210> 26

<211> 2893

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 398269CB1

<400> 26

agtggctgag	tcgggggcgg	gctgggaggg	ctgtcgggtg	gccagttctgc	gtacgacggc	60
ccgtccctctg	cgcacggacg	ccgggaagaa	gggggtgggg	ccacgtttgc	gtccgcgcca	120
tcaggcccgga	gatagcggcg	aggtccgctt	tcagtgtatg	gttttccctg	ccaaacgggt	180
ctgcttggtg	ccatccatgg	agggcggtgcg	ctgggccttt	tcttgccgca	cttggtgcc	240
gagccgagcc	gaatggctgc	tggcagtgcg	atcgattcag	cccaggaga	aggagcgc	300
tggccagttc	gtctttgccc	gggacgctaa	ggcagccatg	gctggctcgc	tgatgataag	360
gaaattagtt	gcagagaaat	tgaatatccc	ttggaatcat	attcgtttgc	aaagaactgc	420
aaaaggaaaa	ccagttcttg	caaaggactc	atcgaatcct	taccggaatt	tcaactttaa	480
catctctcat	caaggagact	atgcagtgc	tgtctgtgaa	cctgagctgc	aagttggaat	540
tgatataatg	aagactagtt	ttccaggtcg	tggttcaatt	ccagaattct	ttcatattat	600
gaaaagaaag	tttaccaca	aagaatggga	aacaatcaga	agctttaagg	atgagtggac	660
tcagctggat	atgttttata	ggaattgggc	acttaaggaa	agcttcataa	aagccattgg	720
tgttggaacta	ggatttgaat	tgcagcggt	tgaatttgat	ctatctccat	taaacttgga	780
tataggccaa	gtttataaag	aaacacgttt	attcctggat	ggagaggaag	aaaaagaatg	840
ggcatttgag	gaaagcaaaa	tagatgagca	ccattttggt	gcagttgctc	ttaggaaacc	900
cgatggatct	agacatcagg	atgttccatc	tcaggatgat	tccaaaccaa	cccagaggca	960
atttactatt	ctcaacttta	atgatttaat	gtcatctgcc	gttcccatga	cacctgaaga	1020
tccttcatttt	tgggactggt	tttgcttcac	agaagaaatt	ccaatacgaa	atgggtacaaa	1080
gtcatgatga	ttccctgagt	aacaaaggga	aatgaaaact	gtttgtgatc	ttccgtattc	1140
actgaaaaat	aaatgcttgt	ttagtatcaa	attttatttc	acgaaagttt	ttttaaagaa	1200
cagaaacttt	tccaattaaa	aaaaaaaagc	agacttctgg	ttcaagatag	ctcactggaa	1260
tacatgttta	cctctttctt	tcctaaattg	cattgaattg	ataggaagga	tggcggaatc	1320
ttaaagtgat	acatgctaac	tgtagaaaaa	aatagaaaat	gcacataagc	aaaagggaac	1380
atttaaatgc	tatctttcaa	agataactac	tcttaaaacc	ttgagtatct	tttcagacct	1440
ttttcttggc	aaatgaatcc	atattgacat	atttgatttt	tttaaaaaca	tggaaacgta	1500

ctgttttgt	aattcttttt	aactgcacat	ctactgttca	taaatatacc	tctgtaacat	1560
aactttttgt	ggttctaagt	tactctgttg	tatagctaata	acaagttagg	atgcttttgg	1620
ccagaggtaa	cagtgtccaa	atataattgg	cctaagtaac	ctaggaaatt	gtttgacata	1680
acacagggtt	caggggtgtc	attaaagaca	cacttttttt	gccttgacct	cagttggttt	1740
gttttgcctt	aggctattcc	acctctcgat	cacaagagct	gctgtataaa	cttccagttt	1800
tacatctttg	taaaattgta	tctcaaaggc	agaaaaggcc	atttcgtctc	tcatttggtt	1860
tatccatgag	gaagattttt	aacaaaagcc	tccagaagat	ttccccctcag	tttccattga	1920
cttagatcag	gttacagaga	aaggcaatgt	ctgacatttt	tggtctctgt	tagaagtaga	1980
ctctgttgaa	aagaaagaag	ctaagctagg	tgtgaagaat	ggaattggaa	gcccactgcc	2040
ttcccataag	aaagggtttac	cataattttac	tcactttttt	ctgtgttaga	catttttgatt	2100
atctgcagtt	tattactaca	agcagtggca	gagtgaatgt	ccttgtagat	tttgagttac	2160
atgcttaatt	atgtccttga	gaaagtttct	aaaagtggaa	tgattgggtt	gactggttca	2220
tagggcttta	attatacaat	ttaccctctc	aattagtact	atatgtatgt	gacttccctc	2280
cccctgccag	aatactcctt	ggtcaattgt	aggatttctt	tttgggttaa	tttttgccaa	2340
tgtaattaaa	aaatgggtatg	tcatttttaa	aatttgtatt	tctttcatta	caaataagat	2400
tgttatgtca	gtattgttat	tggtttttcg	tattcctctt	aacgtgaacc	gtctgttcat	2460
tgtttttacc	tgtttttcgtt	ttagcaagta	ctacttaatt	taaagtgtga	acttaataata	2520
taagatgcca	ggaccatcat	attgatgaca	aaaatctatt	atgtgggtgt	agtgcattgc	2580
ttggagttaa	actcttaagt	atcaagaagt	aaaataattt	cattgtttct	gttactaaca	2640
ttaaaagtgt	gcaaattgta	taaaattagg	tacttcgatt	agtaaaaagaa	aaaaatttga	2700
atgttttttt	ttttatttta	ttaagcatgc	ccagttatgc	caagcatagt	aaataaaggt	2760
caagtagcat	ttataataga	ggaagtattg	ttatccctag	catgagtgtg	atgggtgatg	2820
gaaaaacttt	gtcttgtcat	tataataata	aaaaaatgaa	cattttattat	ggaatttcaa	2880
aaaaaaaaaa	aaa					2893

<210> 27

<211> 2276

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1258888CB1

<400> 27

gagagtgagg	cggaggacc	gagcggctga	ggagagagga	ggcggcggct	tagctgctac	60
gggggtccgg	cggcgccctc	ccgagggggg	ctcaggagga	ggaaggagga	cccgtgcgag	120
aatgcctctg	ccctggagcc	ttgcgctccc	gctgctgctc	tcctgggtgg	caggtgggtt	180
cgggaacgcg	gccagtgcac	ggcatcacgg	gttggttagca	tcggcacgct	agcctggggg	240
ctgtcactat	ggaactaaac	tggcctgctg	ctacggctgg	agaagaaaca	gcaagggagt	300
ctgtgaagct	acatgcgaac	ctggatgtaa	gtttgggtgag	tgctggtggc	caaacaaatg	360
cagatgcttt	ccaggataca	ccgggaaaac	ctgcagtcaa	gatgtgaatg	agtgtggaat	420
gaaacccccg	ccatgccaac	acagatgtgt	gaatacacac	ggaagctaca	agtgtctttg	480
cctcagtggc	cacatgctca	tgccagatgc	tacgtgtgtg	aactctagga	catgtgccat	540
gataaaactgt	cagtacagct	gtgaagacac	agaagaaggg	ccacagtgcc	tggtgccatc	600
ctcaggactc	cgcctggccc	caaatggaag	agactgtcta	gatattgatg	aatgtgcctc	660
tggtaaagtc	atctgtccct	acaatcgaag	atgtgtgaac	acatttggaa	gctactactg	720
caaatgtcac	attggtttcg	aactgcaata	tatcagtggg	cgatatgact	gtatagatat	780
aatgaatgt	actatggata	gccatacgtg	cagccaccat	gccaattgct	tcaataccca	840
agggctcctt	aagtgtaaat	gcaagcaggg	atataaaggc	aatggacttc	ggtgttctgc	900
tatccctgaa	aattctgtga	aggaagtcc	cagagcacct	ggtaccatca	aagacagaat	960
caagaagtgt	cttgctcaca	aaaacagcat	gaaaaagaag	gcaaaaatta	aaaatgttac	1020
cccagaaccc	accaggactc	ctacccttaa	ggtgaacttg	cagcccttca	actatgaaga	1080
gatagtttcc	agaggcgagg	actctcatgg	aggtaaaaaa	gggaatgaag	agaaaatgaa	1140
agaggggctt	gaggatgaga	aaagagaaga	gaaagccctg	aagaatgaca	tagaggagcg	1200
aagcctgcga	ggagatgtgt	ttttccctaa	ggtgaatgaa	gcagggtgaat	tcggcctgat	1260
tctggtccaa	aggaaagcgc	taacttccaa	actggaacat	aaagcagatt	taaatatctc	1320
ggttgactgc	agcttcaatc	atgggatctg	tgactggaaa	caggatagag	aagatgattt	1380
tgactggaa	cctgctgatc	gagataatgc	tattggcttc	tatatggcag	ttccggcctt	1440
ggcagggtca	aagaaagaca	ttggccgagt	gaaactttct	ctacctgacc	tgcaacccca	1500
aagcaacttc	tgtttgcctc	ttgattaccg	gctggccgga	gacaaagtgc	ggaaacttcg	1560
agtgtttgtg	aaaaacagta	acaatgccct	ggcatgggag	aagaccacga	gtgaggatga	1620
aaagtgggaag	acaggggaaaa	ttcagtgtgt	tcaaggaact	gatgctacca	aaagcatcat	1680
ttttgaagca	gaacgtggca	agggcaaaaac	cggcgaaatc	gcagtggatg	gcgtcttgct	1740
tgtttcaggc	ttatgtccag	atagcctttt	atctgtggat	gactgaatgt	tactatcttt	1800
atatttgact	ttgtatgtca	gttccctggg	ttttttgata	ttgcatcata	ggacctctgg	1860
catttttagaa	ttactagctg	aaaaattgtg	atgtaccaac	agaaatatta	ttgtaagatg	1920

```

cctttcttgtg ataagatatg ccaatatattg ctttaaatat catatcactg tatctttctca 1980
gtcattttctg aatcctttcca cattatatatta taaaatatgg aaatgtcagt ttatctcccc 2040
tcctcagtat atctgatttg tataagtaag ttgatgagct tctctctaca acattttctag 2100
aaaatagaaa aaaaagcaca gagaaatgtt taactgtttg actcttatga tactttcttg 2160
aaactatgac atcaaagata gacttttgcc taagtggctt agctgggtct ttcatagcca 2220
aacttgata tttaaattct ttgtaataat aatatccaaa tcatcaaaaa aaaaaa 2276

```

```

<210> 28
<211> 2016
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1375891CB1

```

```

<400> 28
gaaaagggtac ccgcgagaga cagccagcag ttctgtggag cagcgggtggc cggctaggat 60
gggctgtctc tgggggtctgg ctctgcccc tttcttcttc tgctgggagg ttgggggtctc 120
tgggagctct gcaggcccca gcaccgcag agcagacact gcgatgacaa cggacgacac 180
agaagtgtcc gctatgactc tagcacggg ccacgcgct ctggaaactc aaacgtgag 240
cgctgagacc tcttctaggg cctcaacccc agccggcccc attccagaag cagagaccag 300
gggagccaag agaatttccc ctgcaagaga gaccaggagt ttcacaaaaa catctcccaa 360
cttcatgggtg ctgatcgcca cctccgtgga gacatcagcc gccagtggca gccccgaggg 420
agctggaatg accacagttc agaccatcac aggcagtgat cccgaggaag ccatctttga 480
caccctttgc accgatgaca gctctgaaga ggcaaagaca ctcaaatgg acatattgac 540
attggtcac acctccacag aagctaaggg cctgtcctca gagagcagtg cctcttccga 600
cggcccccat ccagtcatca ccccgtcacg ggcctcagag agcagcgcct ctcccgacgg 660
cccccatcca gtcatcaccc cgtcacgggc ctgagagagc agcgccctct ccgacggccc 720
ccatccagtc atcaccccgt catggtcccc gggatctgat gtcactctcc tegtgaagc 780
cctggtgact gtcacaaaaca tcgaggttat taattgcagc atcacagaaa tagaaacaac 840
aacttccagc atccctgggg cctcagacat agatctcatc cccacggaag ggggtgaaggc 900
ctcgtccacc tccgatccac cagctctgcc tgactccact gaagcaaaac cacacatcac 960
tgaggtcaca gcctctgccg agaccctgtc cacagccggc accacagagt cagctgcacc 1020
tcattgccacg gttgggaccc cactccccac taacagcgcc acagaaagag aagtgcacgc 1080
accgggggac acgaccccca gtggagctct ggtcacagtt agcaggaatc cctggaaga 1140
aacctcagcc ctctctgttg agacaccaag ttacgtcaaa gtctcaggag cagctccggg 1200
ctccatagag gctgggtcag cagtgggcaa aacaacttcc tttgctggga gctctgcttc 1260
ctcctacagc cctcgggaag ccgcccctca gaacttcacc ccttcagaga caccgacctc 1320
ggacatcgca accaaggggc cctccccac cagcagggac cctcttccct ctgtccctcc 1380
gactacaacc aacagcagcc gagggacgaa cagcacctta gccaatgca caacctcagc 1440
gaagaccacg atgaagcccc aacagccacg cccacgactg cccggacgag gccgaccaca 1500
gacgtgagtg caggtgaaaa tggaggtttc ctccctctgc ggctgagtg ggcttccccg 1560
gaagacctca ctgaccccag agtggcagaa aggtctgatgc agcagctcca cggggaactc 1620
cacgccacg cgctcactt ccaggtctcc ttactgcgtg tcaggagagg taacgggaca 1680
tcagctgcag ccaggcatgt cccgtatgcc aaaagagggt gctgccccta gcctggggcc 1740
ccaccgacag actgcagctg cgttactgtg ctgagaggta cccagaaggt tcccatgaag 1800
ggcagcatgt ccaagcccc gaccccagat gtggcaacag gaccctcgct cacatccacc 1860
ggagtgtatg tgtggggagg ggcttcacct gttcccagag gtgtccttgg actcaccttg 1920
gcacatgttc tgtgtttcag taaagagaga cctgatcacc catctgtgtg ctccatect 1980
gcattaaaaat tcactcagtg tggcccaaaa aaaaaa 2016

```

```

<210> 29
<211> 2520
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1524355CB1

```

```

<400> 29
caagatggcg gctggcggag ctgtcgctgc ggcgcccag tgccggcttc tcccctacgc 60
gtacacaaag tggagctcct tttcctccac ctacctccc gagaacattt tagtggacaa 120
accaaagac caatcttcaa gatggtcttc agagagcaac tatcctccc agtacttgat 180
tctaaagctc gaaaggcctg ctatagtcca gaatatcaca tttggaaaat atgagaaaac 240
tcatgtttgc aatttgaaga aatttaaagt ctttggtgga atgaatgaag aaaatatgac 300

```

agagctgttg	tccagtggct	taaagaatga	ttataacaaa	gaaacattca	ccttgaagca	360
taaaattgat	gaacagatgt	tcccttgtcg	attcattaaa	atagttccac	tcttgtcctg	420
gggacccagc	tttaacttta	gcattctggt	tgttgaactt	agtggcattg	atgatacctga	480
tatagtacaa	ccttgtctca	actggtatag	caagtaccgt	gaacaggaag	ctattcgcct	540
ttgcctaaaa	cacttcagac	aacacaacta	tacagaagct	tttgagtcac	tgcaaaagaa	600
aaccaagatt	gcactggaac	atccccatgtt	aacagatatt	catgacaagc	tgggtgtttaa	660
gggtgatttt	gatgcttgcg	aagagttgat	tgaaaaggct	gtaaatgatg	gcttgttcaa	720
tcagtatatc	agtcaacagg	aatataagcc	acgatggagt	caaatacattc	ccaaaagtac	780
caaaggtgat	ggggaagata	accgtccagg	aatgagagga	ggccatcaga	tggttattga	840
tgttcaaaca	gagactgttt	atttgtttgg	tggctgggat	ggaacacaag	atcttgcctga	900
cttctgggcg	tacagtgtga	aggagaacca	gtggacatgt	atctctagag	acactgaaaa	960
agagaattgt	cctagtgtcca	gatcgtgtca	taaaatgtgc	attgatattc	aacggaggca	1020
aatctacaca	ttggggcggt	acttggattc	ctctgtgagg	aacagcaaat	ctctgaaaag	1080
tgacttctat	cgttatgaca	ttgatacaaa	cacatggatg	ttactaagtg	aggatactgc	1140
tgctgatgga	gggccgaaat	tgggtgttga	tcatacagatg	tgtatggact	cagaaaaaca	1200
tatgatctac	acttttgggt	gtagaatttt	gacttgtaat	ggcagcgtag	atgacagcag	1260
agccagtga	ccacaattca	gtggctgttt	tgctttcaac	tgtcaatgtc	aaacctggaa	1320
acttcttcga	gaggactcct	gtaatgctgg	gcctgaggac	atccagtcctc	gaataggaca	1380
ctgcatgtta	ttccactcaa	aaaatcgttg	cttatatgta	tttgggtggcc	agcgatcaaa	1440
gacctatttg	aatgatttct	ttagtattga	tgtggactct	gatcatgtag	acataatatc	1500
agattggcacc	aagaaagact	ctgggatggg	tccaatgaca	ggattttacac	agagagcaac	1560
tattgatcca	gaactgaatg	aaatacacgt	cttatctgga	ctcagcaaa	ataaggaaaa	1620
gagggaagaa	aatgttagaa	attcattctg	gatttatgac	attgtgagga	atagttgggtc	1680
ttgtgtctat	aagaatgatc	aagctgcaaa	ggataatcca	actaaaagtc	ttcaggaaga	1740
agaacctagt	ccaaggtttg	cccatacagc	tgtatacgat	gagctacaca	aggttcatta	1800
cttatttggg	gggaatccag	gaaaatcttg	ctctccaaag	atgagattag	atgacttctg	1860
gtcactgaag	ttgtgtagac	cttcaaaaga	ttatttactg	aggcattgca	agtaacctcat	1920
aagaaaacac	aggtttgaag	aaaaggccca	agtggatccc	cttagtgctc	tgaaatattt	1980
acaaaatgat	ctttatataa	ctgtggatca	ttcagaccca	gaagagacaa	aagagtttca	2040
gctcttgga	tcagctctat	tcaaatcttg	ttcagatttt	acagctctgg	gcttttctga	2100
tgtggatcac	acctatgctc	aaagaactca	gctctttgac	accttagtaa	atttttctcc	2160
tgacagcatg	actcctccta	aaggcaacct	ggtagacctc	atcacactgt	aactgaagag	2220
tcactggaca	cagaaatgga	aaacaggagt	cgattttccg	tcttttggat	tgcagctcca	2280
ctgactgaca	gtaaagctgc	agtgattgag	gactgcacca	gagttctgaa	gggatcttaa	2340
ccatcacaa	tttttaccct	cttccctcat	gcttgacctc	aaccccgcctc	tctctatcct	2400
attcctaaat	taggtgaaata	aagtgaattt	ggtatacttt	ccagttaaat	atatatatat	2460
atatattttt	tcttacttta	tcttttaaga	attaatgagt	ataaaagcaa	aattaggcag	2520

<210> 30

<211> 1954

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1598937CB1

<400> 30

ccgagttatg	cagctccggg	cggaagggg	tcttctcgag	gggcgcggtc	cataagggtt	60
gaggccagag	gcgcgtacct	ctgggcgcgg	agctctggag	tggaggctgg	ggttcggagt	120
gcgtcatctg	gaagcaggca	ccccggccgg	caggcagagt	cacgggtggca	gcattgagag	180
ttggacaccc	gggtccttga	agtgatctct	aggccccagc	cccaaatccg	ccaccattcc	240
gtgctgcggg	gacaccatgg	ctccagaaga	ggacgctgga	ggggaggcct	tagggggcag	300
tttctgggag	gctggcaact	acaggcgcac	ggtacagcgg	gtggaggacg	ggcaccggct	360
gtgcggggac	ctggtcagct	gcttccagga	gcgcgcgcgc	atcgagaagg	cttatgcccc	420
gcagttggct	gactgggccc	gaaagtggag	ggggaccgtg	gagaaggggc	cccagtatgg	480
cacactggag	aaggcctggc	atgccttttt	cacggcggtc	gagcggctga	gcgcgctgca	540
cctggagggtg	cgggagaagc	tgcaagggca	ggacagttag	cgggtgcgcg	cctggcagcg	600
gggggcttct	caccggcctg	tgctgggcgg	cttccgcgag	agccgggcgg	ccgaggacgg	660
cttccgcaag	cccagaagc	cctggctgaa	gaggctgaag	gaggttgagg	cttccaagaa	720
aagctaccac	gcagcccgga	aggatgagaa	gaccgcccag	acgaggggaga	gccacgcaaa	780
ggcagacagc	gccgtctccc	aggagcagct	gcgcaactg	caggaacggg	tggaaacgtg	840
tgccaaggag	gccgagaaga	caaaagctca	gtatgagcag	acgctggcag	agctgcacg	900
ctacactcca	cgctacatgg	aggacatgga	acaggccttt	gagacctgcc	aggccgcccga	960
gcgcacggcg	cttcttttct	tcaaggatat	ctgtctcacc	ttacaccagc	acctggacct	1020
ttccagcagt	gagaagttcc	atgaactcca	ccgtgacttg	caccagggca	ttgaggcagc	1080
cagtgcagaa	gaggatctgc	gctgggtggcg	cagcaccac	gggccaggca	tggccatgaa	1140

ctggccacag	tctgaggagt	ggtccttggg	cacacagagg	acaatcagcc	ggaaagagaa	1200
gggtggccgg	agccctgatg	aggttaccct	gaccagcatt	gtgcctacaa	gagatggcac	1260
cgcaccccc	ccccagtcct	cggggctccc	aggcacgggg	caggatgagg	agtgggtcaga	1320
tgaagagagt	ccccggaagg	ctgccaccgg	ggttcgggtg	agggcactct	atgactacgc	1380
tggccaggaa	gctgatgagc	tgagcttccg	agcaggggag	gagctgctga	agatgagtga	1440
ggaggacgag	cagggctggg	gccaaggcca	gttgcagagt	ggccgcattg	gcctgtaccc	1500
tgccaactac	gtggagtgtg	tgggcgcctg	agtgtcctga	cagcccttct	gcaacgttta	1560
cccaccctgg	tctcagagccc	agcttctcct	ggagagccgg	accctcaggg	ccctgaaccg	1620
tgcctctctg	gctgctcctc	tgtcccttga	gggaggaagt	cctgggaccc	agggagggga	1680
ggggcctttg	tctaggggaag	ggactggtag	ggaagggacg	agtctaggct	gagggcaaga	1740
tgggagggtca	gaggtgacag	aagcgttcag	gggtgcctgg	gcctccccag	gagctgtgga	1800
ctcagttcct	gacctctgct	ttggggttcc	tgggggtggc	ttgggggtgag	tgtagttctg	1860
gcctagcagc	accctcttgt	ggcttgttct	agcgtgtatt	aaaacttgac	acacaccac	1920
acacaaaaac	aaaaacacca	aaaaaaaaaa	aaaa			1954

<210> 31
 <211> 1817
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1725801CB1

<400> 31

gacgcggtga	ggagacggcc	cacggcgccc	gcgggctggg	gcggctcgtt	cttccttctc	60
cgtggcctac	gagggctccc	agcctgggta	aagatggccc	catggccccc	gaagggccta	120
gtcccagctg	tgctctgggg	cctcagcctc	ttcctcaacc	tcccaggacc	tatctggctc	180
cagccctctc	cacctcccca	gtcttctccc	ccgcctcagc	cccatccgtg	tcatacctgc	240
cggggactgg	ttgacagctt	taacaagggc	ctggagagaa	ccatccggga	caacttttga	300
ggtggaaaca	ctgcctggga	ggaagagaat	ttgtccaaat	acaaagacag	tgagaccgcc	360
ctggtagagg	tgctgggggg	tgtgtgcagc	aagtcagact	tcgagtgcc	ccgcctgctg	420
gagctgagtg	aggagctggg	ggagagctgg	tggtttcaca	agcagcagga	ggccccggac	480
ctcttccagt	ggctgtgtct	agattccctg	aagctctgct	gccccgcagg	caccttcggg	540
ccctcctgcc	ttccctgtcc	tgggggaaca	gagaggccct	gcgggtggcta	cgggcagtg	600
gaaggagaag	ggacacgagg	gggcagcggg	cactgtgact	gccaagccgg	ctacgggggt	660
gagggccttg	gccagtgtgg	ccttggctac	tttgaggcag	aacgcaacgc	cagccatctg	720
gtatgttcgg	cttgttttgg	cccctgtgcc	cgatgtctag	gacctgagga	atcaaactgt	780
ttgcaatgca	agaagggctg	ggccctgcat	cacctcaagt	gtgtagacat	tgatgagtgt	840
ggcacagagg	gagccaactg	tggagctgac	caattctgcg	tgaacactga	gggtccttat	900
gagtgcggag	actgtgccaa	ggcctgccta	ggctgcatgg	gggcagggcc	aggtcgctgt	960
aagaagtgtg	gccttggtta	tcagcagggt	ggctccaagt	gtctcgatgt	ggatgagtgt	1020
gagacagagg	tgtgtccggg	agagaacaag	cagtgtgaaa	acaccgaggg	cggttatcgc	1080
tgcatctgtg	ccgaggggcta	caagcagatg	gaaggcatct	gtgtgaagga	gcagatccca	1140
gagtcagcag	gcttcttctc	agagatgaca	gaagacgagt	tggtgggtgct	gcagcagatg	1200
ttctttggca	tcatacatctg	tgactgggcc	acgctggctg	ctaagggcga	cttgggtgtc	1260
accgccatct	tcattggggc	tgtggcgggc	atgactggct	actgggtgtc	agagcgcagt	1320
gaccgtgtgc	tggagggctt	catcaagggc	agataatcgc	ggccaccacc	tgtaggacct	1380
cctcccaccc	acgctgcccc	cagagcttgg	gctgccctcc	tgctggacac	tcaggacagc	1440
ttggtttatt	tttgagagtg	gggtaagcac	ccctacttgc	cttacagagc	agcccaggtg	1500
cccaggcccc	ggcagacaag	gcccctgggg	taaaagttag	ccctgaaggt	ggataccatg	1560
agctcttcac	ctggcggggg	ctggcaggct	tcacaatgtg	tgaatttcaa	aagtttttcc	1620
ttaatgggtg	ctgctagagc	tttggccctt	gcttaggatt	aggtgggtcct	cacaggggtg	1680
gggccatcac	agctccctcc	tgccagctgc	atgctgccag	ttcctgttct	gtgttcacca	1740
catccccaca	ccccattgcc	acttatttat	tcattctcagg	aaataaagaa	aggtcttgga	1800
aagttaaaaa	aaaaaaa					1817

<210> 32
 <211> 2694
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1730482CB1

<400> 32

gacctagtgt	gagcataatg	gaaaaaacac	aatcacttcc	tacacgacca	ccaacttttc	60
ctccaacccat	tccaccagca	aaagaagtat	gtaaggcggc	caaggctgac	ctgggtattta	120
tgggtggatgg	atcctggagc	attggagatg	aaaatttcaa	taagatcatc	agcttttctat	180
acagcactgt	tggagccctg	aacaagattg	gcacagatgg	aaccaagtt	gcaatgggttc	240
agttcactga	tgatcccaga	acagaattta	aactaaatgc	ttacaaaacc	aaagagactc	300
ttcttgatgc	aattaaacac	atttcataca	aaggaggaaa	tacaaaaaca	ggaaaagcaa	360
ttaagtatgt	tcgagatacc	ttgttcactg	cagagtcagg	tacaagaagg	ggcatcccaa	420
agggttatcgt	ggttataact	gatggaagat	cacaagatga	tgtgaacaaa	atctccaggg	480
agatgcaatt	agatggctat	agcatttttg	caattgggtg	ggccgatgca	gattactcgg	540
agttgggttag	cattggcagt	aagcccagcg	cacgccatgt	cttctttgtg	gatgactttg	600
acgccttttaa	gaaaatcgaa	gatgagttaa	ttacttttgt	ctgcgaaaca	gcatacagcaa	660
cctgtccagt	ggtacacaag	gatggcattg	atcttgcagg	atttaagatg	atggaaatgt	720
ttggtttggg	tgaaaaagat	ttttcatcag	tgggaagggg	ttctatggag	cctgggtacct	780
tcaatgtgtt	tccatgttac	caactccata	aagatgccct	ggtttcccag	ccaaccagggt	840
acttgcaccc	agaaggattg	ccctccgact	acacaatcag	ttttctattc	cggatttctc	900
ctgacactcc	acaggagcca	tttgcctttt	gggagatttt	aaataaaaaat	tctgacctat	960
tggttggggg	tatttttagac	aatgggtggga	aaactctaac	atatttcaac	tatgcaccaga	1020
gtgggggattt	tcaaactggt	actttcgaag	gacctgaaat	taggaaaatt	ttttatggaa	1080
gctttcacaa	gctacacatt	gttgtcagtg	agactttggg	caaagtgggt	attgactgca	1140
agcaagtggg	tgagaaggca	atgaacgcac	cagctaatat	cacgtcagat	ggtgtagaag	1200
tgctagggga	aatgggtcga	tcaagaggac	cagggtgaaa	ctctgcaccg	ttccagttac	1260
agatgtttga	tattgtttgc	tccacatcat	gggccaatac	agacaaatgc	tgtgaacttc	1320
caggcctgag	agatgatgag	tcttgcccag	accttcccca	ttcctgctcc	tgttctgaaa	1380
ccaatgaagt	ggctctggga	ccagcggggc	caccagggtg	tccaggactc	cgaggaccaa	1440
agggccagca	aggtgaaccg	ggtccaaagg	gaccagatgg	ccctcggggg	gaaattggtc	1500
tgccaggacc	tcagggtcca	cctggacctc	aaggaccaag	tggctctgtc	attcaaggaa	1560
tgcccggaat	gccaggagaa	aaaggagaga	aaggagatac	tggccttcca	ggtccacagg	1620
gtatcccagg	aggcgttggt	tcaccaggac	gtgatggctc	accaggccag	aggggccttc	1680
cgggaaagga	tggatcctcg	ggacctccag	gaccaccagg	gccaataggc	attccttgca	1740
ccctctggag	cccagggatc	acaggaagca	tgggaccgca	aggcgccctg	ggaccacctg	1800
gtgtccctgg	agcaaaaggg	gaacgaggag	agcggggtga	cctgcagtct	caagccatgg	1860
tgagatcagt	ggcgcgctcaa	gtatgcgaac	agctcatcca	gagtcacatg	gccagggtaca	1920
ctgccatcct	caaccagatt	cccagccact	ctcatccat	ccggactgtc	caagggcctc	1980
ctggggagcc	tgggaggcca	ggctcacctg	gagcccttgg	tgaacaagga	cccccaggca	2040
caccaggctt	ccccggaaat	gcaggcgtgc	cagggaaccc	aggagaacga	ggtctaactg	2100
gtatcaaagg	agaaaaagga	aatccaggcg	ttggaaccca	aggtccaaga	ggccccctg	2160
gaccagcagg	accttcaggg	gagagtcggc	ctggcagccc	tgggccccct	ggctctcctg	2220
gaccaagagg	cccaccagg	catctggggg	ttcctggacc	ccaaggctct	tctggccagc	2280
ctggatatgt	tgacccctca	tcatgttctg	cctatgggtg	gagagctccc	catccagatc	2340
agccagagtt	cacccctgtc	caagatgagc	tgggaagccat	ggaactgtgg	ggccctggag	2400
tctgatagcc	tcaggagaaa	tttgaagacc	aactgcaaga	actcttaagg	aatcttggtt	2460
gagaaaatgt	tgttatgtgg	tttgtatgct	acttttgggg	ggcagggtct	atttcagcag	2520
cctaaatctc	ctccttggtg	aatgttaata	ttattattat	tattaacaaa	aaatatatat	2580
ttttaaaaag	ttcccttaat	ctatgacatg	gtagcaatga	tttccctttg	gtgtcttaat	2640
ggcatgtcag	ataatttggt	tttccagaga	agagagctca	aagaggaatt	ggga	2694

<210> 33

<211> 1149

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1810058CB1

<400> 33

cagtatctgg	gtccagcctg	cagccttagg	gtccagggtga	tgttaccgtg	tgtgtggccc	60
ttcttcacag	tggcctccta	gaaaaacaag	accctgactc	aaagaacacc	tctcactaca	120
ttcagagtct	gtcatctgaa	ccatgaggat	ctgggtggctt	ctgcttgcca	ttgaaatctg	180
cacagggaac	aggacacctg	ataaactcac	caggcaaggg	cacctggaa	ttcctgggaa	240
ccccggtcac	aatgggtctg	ctggaagaga	tggacgagac	ggagcgaagg	gtgacaaagg	300
cgatgcagga	gaaccaggac	gtcctggcag	cccggggaag	gatgggacga	gtggagagaa	360
gggagaacga	ggagcagatg	gaaaagttag	agcaaaaggc	atcaaagggtg	atcaaggctc	420
aagaggatcc	ccaggaaaaa	atggccccc	ggggcttgca	gggcccattg	gagagaaagg	480
cctccgagga	gagactgggc	ctcaggggca	gaagggaat	aagggtgacg	tgggtcccac	540
tggctcctgag	gggccaaggg	gcaacattgg	gcctttgggc	ccaactgggt	taccgggccc	600
catgggccc	attggaaagc	ctgggtccca	gggagaagct	ggaccacagg	ggccccaggg	660

tgagccagga	gtccggggaa	taagaggctg	gaaaggagat	cgaggagaga	aagggaaaaat	720
cgggtgagact	ctagtcttgc	caaaaagtg	tttctactgtg	gggctcacgg	tgctgagcaa	780
gtttccttct	tcagatgtgc	ccattaaatt	tgataagatc	cacatcactg	ttttctccag	840
gaatgttcag	gtgtctttgg	tcaaaaacgg	agtaaaaata	ctgcacacca	gagatgctta	900
cgtgagctct	gaggaccagg	cctctggcag	cattgtcctg	cagctgaagc	tccgggatga	960
gatgtggctg	cagggtgacg	gaggagagag	gttcaatggc	ttgtttgctg	atgaggacga	1020
tgacacaact	ttcacagggt	tccttctgtt	cagcagccag	tgacagagga	gagtttataa	1080
atctgccaga	ccatccatca	gaatcagctt	gggatgaact	tattcagatg	gttttacttt	1140
attaattca						1149

<210> 34

<211> 1215

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2040679CB1

<400> 34

gaagaactag	catgtatgta	ttatctccag	tggaatttat	aattctacaa	cttttatttta	60
ttcaggccat	ttccagcagt	ttaaaagggt	tcctttcagc	tatgagactg	gctcatagag	120
gctgtaatgt	tgatacacca	gtttcaacgc	tcacaccagt	gaagacttca	gaatttgaaa	180
actttaaaac	taaaatgggt	atcacatcca	aaaaagacta	tcctctaagt	aagaattttc	240
catattcctt	ggaacatctt	cagacttctt	actgtgggct	tgcccgagtt	gatatgcgta	300
tgcttttgctt	aaaaagcctt	aggaatagg	acttgagtca	caaccatata	aaaaagcttc	360
cagctacaat	tgagacctc	atacaccttc	aagaacttaa	cctgaatgac	aatcacttgg	420
agtcatttag	tgtagccttg	tgtcattcta	cactccagaa	gtcacttcgg	agtttggacc	480
tcagcaagaa	caaaatcaag	gcactccctg	tcgagttttg	ccagctccag	gaacttaaga	540
atttaaaact	tgacgataat	gaattgattc	aatttccttg	caagatagga	caactaataa	600
accttcgctt	tttgtcagca	gctcgaaata	agcttcattt	tttgccctagt	gaatttagaa	660
atztatccct	tgaatacttg	gatctttttg	gaaatacttt	tgaacaacca	aaagtccttc	720
cagtaataaaa	gctgcaagca	ccattaaact	tattggaatc	ttctgcacga	accatattac	780
ataataggat	tccatattgg	tctcatatca	ttccattcca	tctctgccaa	gatttggata	840
ccgcaaaaat	ttgtgtttgt	ggaagattct	gtctgaactc	tttcattcaa	ggaactacta	900
cctgaatct	gcattctgtt	gcccacactg	tggtcttagt	agataatttg	ggtggtagctg	960
aagcacctat	tatctcttat	ttctgttctc	taggctgtta	tgtaatttcc	tctgatattg	1020
taaagtaatg	ggtgagacca	gaaaaagaaa	tttcaataac	agatcagttt	ggggtgcatg	1080
tatgattttg	cagcgtcaaa	ttggagtaag	ggaagatttc	tgtatacttg	ctggagagga	1140
ggaatgtgta	tagttactca	tttagatgac	tccaaaactt	ttattaaaac	caatttttagt	1200
tttaaaaaaa	aaaaa					1215

<210> 35

<211> 1300

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2960051CB1

<400> 35

ttctgcacaa	agaactgaaa	ggcattttatc	cccaagggag	gcagttattt	tagatttttac	60
taagaagttc	agcaaatact	tttcaacatt	cccttctgtc	ctttctttgt	ttttaagaa	120
agctctgatt	ttgttttcatt	ttcagctgga	gactttaaag	acaccaagca	aagcctactt	180
agtttagatc	tccagaaaatt	ggctgggtgga	aaaaaatcaa	acatgaagat	tgagtttttg	240
ttttgttttt	ttctgcttat	cattttttcaa	actgactttg	gaaaaaatga	agaaattcct	300
aggaagcaaa	ggaggaagat	ctaccacaga	aggttgagga	aaagttcaac	ctcacacaag	360
cacagatcaa	acagacagct	tgggaattccg	caaacaacag	tttttacacc	agtagcaaga	420
cttctatttg	tttaactttga	ttatagcatg	gaggaaaagt	ttgaatcctt	ttcaagtttt	480
cctggagtag	aatcaagcta	taatgtgtta	ccaggaaaga	agggacactg	tttggtaaag	540
ggcataacca	tgtacaacaa	agctgtgtgg	tcgcctgagc	cctgcactac	ctgcctctgc	600
tcagatggaa	gagttctttg	tgatgaaacc	atgtgccatc	cccagagggtg	cccccaaaca	660
gttatacctg	aaggggaatg	ctgccctgtc	tgctccgcta	ctgggtacaga	gatttagcta	720
agcaaaatat	cagtgtgtga	ttaatcttta	acttccattt	gtttttgtta	ctaatttttag	780
attaaaatta	tgatacatta	gtcagatctg	agtacttaaa	atattggcaa	aatgctgatt	840
aacatagaaa	atatctggga	aaatgtatgg	taggggatat	aaataataga	ctgtggcttt	900


```

atagtttctag ctctatcaga ttcagtaaac ttggatgaga ttacattcca catttgactc 960
tcagcttttag agatatggta acagaatttc tacaacagat cctgaattct tattgcatta 1020
agggctctgc tttggtctat atgtgcatta tcccacttaa tccagtgcga cgtgccttta 1080
tcaccttgaa gccagggtaa acaaaggaag agtgatttgc atctaaagag aacaaagccc 1140
caaccctctg gctataccca accactcaaa ggcagcacag gaaccacat cactgcttgg 1200
ataatcccag gaaaatgcag aaaaagtgtg gcctgaagca tgattttctc atgtggcact 1260
tctgtgtgca ggagatcaca gcgcggtttt gttgctgcca 1300

```

```

<210> 36
<211> 1562
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 3117318CB1

```

```

<400> 36
aaggccggggc gcggttaagag cgtctcgggg agtagggcaa ggccggccggg cccctcccat 60
tccgcctttt cttcagcgtc ctgcccgcgg cactggctgc ggggtgccggg ccacctgcga 120
gtgtgcgcag ggactctgga caccgcgggc ggcgagctga gggagcagtc tccacgagga 180
cccaggcgga cctctggcg ccatgcgcgc cctcccggc ctgctggagg ccagggcgcg 240
tacgccccgg ctgctcctcc tccagtgcct tctcgtgcc gcgcgccaa gctcggcgga 300
cggcagtgcc ccagattcgg cttttacaag tccacctctc agagaagaaa taatggcaaa 360
taacttttcc ttggagagtc ataacatatc actgactgaa cattctagta tgccagtaga 420
aaaaaatatc actttagaaa ggccttctaa tgtaaattct acatgccagt tcacaacatc 480
tgggggatttg aatgcagtaa atgtgacttg gaaaaaagat ggtgaacaac ttgagaataa 540
ttatcttgtc agtgcaacag gaagcacctt gtatacccaa tacaggttca ccatcattaa 600
tagcaaacaa atgggaagtt attcttgttt ctttcgagag gaaaaggaac aaaggggaac 660
atttaatttc aaagtccctg aacttcatgg gaaaaacaag ccattgatct cttacgtagg 720
ggattctact gtcttgacat gtaaatgtca aaattgtttt ctttaaatt ggacctggtg 780
cagtagtaat gggagtgtaa aggttcctgt tgggtgtcaa atgaataaat atgtgatcaa 840
tggaacatat gctaacgaaa caaagctgaa gataacacaa cttttggagg aagatgggga 900
atcttactgg tgccgtgcac tattccaatt aggcgagagt gaagaacaca ttgagcttgt 960
ggtgctgagc tatttggtgc cctcaaaacc atttcttgta atagtggctg aggtgattct 1020
tttagtggcc accattctgc tttgtgaaaa gtacacacaa aagaaaaaga agcactcaga 1080
tgaggggaaa gaatttgagc agattgaaca gctgaaatca gatgatagca atggatataga 1140
aaataatgtc cccaggcata gaaaaaatga gtctctgggc cagtgaatac aaaacatcat 1200
gtcgagaatc attggaagat atacagagtt cgtatttcag ctttgtttat ccttctctgt 1260
aagagcctct gagtttttag ttttaaaagg atgaaaagct tatgcaacat gctcagcagg 1320
agcttcacat acgatatatg tcagatctaa aggtatatatt tcattctgta attatgttac 1380
ataaaagcaa tgtaaatcag aataaatatg ttagaccaga atacaaatta attatatatt 1440
ggtcttcaaa ggacacacag aacagatatc agcagaatca ctttaactt catagaacaa 1500
aaatcactca aaacctgttt ataaccaaag aattcatgaa aaagaaagcc tttggccatt 1560
tg 1562

```

```

<210> 37
<211> 2801
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> unsure
<222> 2793
<223> a, t, c, g, or other

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 3486992CB1

```

```

<400> 37
gacttttgctt gaatgtttac attttctgct cgctgtccta catatcacaa tatagtgttc 60
acgttttggtt aaaactttgg ggtgtcagga gttgagcttg ctacagcaagc cagcatggct 120
aggatgagct ttgttatagc agcttgccaa ttgggtgctgg gcctactaat gacttcatta 180
accgagtctt ccatacagaa tagtgagtgt ccacaacttt gcgtatgtga aattcgctcc 240
tggtttaccc cacagtcaac ttacagagaa gccaccactg ttgattgcaa tgacctccgc 300
ttaacaagga ttcccagtaa cctctctagt gacacacaag tgcttctctt acagagcaat 360

```

```

aacatcgaga agactgtgga tgagctgcag cagcttttca acttgactga actagatttc 420
tcccaaaaaca actttactaa cattaaaggag gtcgggctgg caaacctaac ccagctcaca 480
acgctgcatt tggaggaaaa tcagattacc gagatgactg attactgtct acaagacctc 540
agcaaccttc aagaactcta catcaaccac aaccaaatta gcactatttc tgctcatgct 600
tttgcaggct taaaaaatct attaaggctc cacctgaact ccaacaaatt gaaagtattt 660
gatagtcgct ggtttgattc tacacccaac ctggaaattc tcatgatcgg agaaaaccct 720
gtgattggaa ttctggatat gaacttcaaa cccctcgcaa atttgagaag cttagttttg 780
gcaggaatgt atctcactga tattcctgga aatgcttttg tgggtctgga tagccttgag 840
agcctgtctt tttatgataa caaactgggt aaagtccctc aacttgccct gcaaaaagtt 900
ccaaatttga aattcttaga cctcaacaaa aaccccatte acaaaatcca agaaggggac 960
ttcaaaaata tgcttcgggt aaaagaactg ggaatcaaca atatgggaga gctcgtttct 1020
gtcgaccgct atgccctgga taacttgccct gaactcaca agctggaagc caccaataac 1080
cctaaactct cttacatcca ccgcttggtt ttccgaagtg tccctgctct ggaaagcttg 1140
atgctgaaca acaatgcctt gaatgccatt taccaaaaga cagtcgaatc cctccccaat 1200
ctgcgtgaga tcagtatcca tagcaatccc ctcagggtgtg actgtgtgat ccactggatt 1260
aactccaaca aaaccaacat ccgcttcatt gagccctgt ccatgttctg tgccatgccg 1320
cccgaaatata aagggcacca ggtgaaggaa gttttaatcc aggattcgag tgaacagtg 1380
ctcccaatga tatctcacga cagcttccca aatcgtttta acgtggatat cggcacgacg 1440
gttttcttag actgtcgagc catggctgag ccagaacctg aaatttactg ggtcactccc 1500
attggaaata agataactgt ggaaccctt tcagataaat acaagctaag tagcgaaggt 1560
accttggaaa tatctaacta acaaattgaa gactcaggaa gatacacatg tgttgcccag 1620
aatgtccaaag gggcagacac tcgggtggca acaattaaag ttaacgggag ccttctggat 1680
ggtacccagg tgctaaaaat atacgtcaag cagacagaat cccattccat cttagtgtcc 1740
tggaaggtta attccaatgt catgacgtca aacttaaaat ggtcgtctgc caccatgaag 1800
attgataacc ctcacataac atatactgcc agggctccag tcgatgtcca tgaatacaac 1860
ctaacgcac cagagccttc cacagattat gaagtgtgtc tcaagtgtc caatattcat 1920
cagcagactc aaaagctatg cgtaaatgtc acaaccaaaa atgccgcctt cgcagtggac 1980
atctctgac aagaaaccag tacagccctt gctgcagtaa tggggtctat gtttgccgtc 2040
attagccttg cgtccattgc tgtgtacttt gccaaaagat ttaagagaaa aaactaccac 2100
cactcattaa aaaagtatat gcaaaaaacc tcttcaatcc cactaaatga gctgtacca 2160
ccactcatta acctctggga aggtgacagc gagaaagaca aagatggttc tgcagacacc 2220
aagccaacc aggtcgacac atccagaagc tattacatgt ggtaactcag aggatatttt 2280
gcttctggta gtaaggagca caaagacgtt tttgctttat tctgcaaaag tgaacaagtt 2340
gaagactttt gtatttttga ctttgctagt ttgtggcaga gtggagagga cgggtggata 2400
tttcaattt ttttagtata gcgtatcgca agggtttgac acggctgcca gcgactctag 2460
gcttccagtg tgtgtttggt ttttattctt ttctatttta tgattgttat tatatgaaa 2520
ttttatttta gttgttgtgc taaactcaat aatgctgttc taactacagt gctcaataaa 2580
atgattaatg acaggatggg gttccctgt gcttttacca gtagcatgac cccttctgaa 2640
gccatccgta gaaagtactt tgtcccaaaa aagcaacata cgggttgaac agcatgaaac 2700
ttttagcat cgggctaaga ctttaactca gagcaaggca gactggtacc tcgttaagat 2760
gtagtactg cggatgttta cactgaatga agntgcttaa t 2801

```

<210> 38

<211> 2597

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4568384CB1

<400> 38

```

ccagaggcaa agaggccagt gaggactgct ctgtgcagtt gtccaggcct gaagaagggtg 60
gcgggtgatt gaatcagaag aggtggcatt ctcttcatta ggaggatatg gataatcaag 120
acctaacaaa ttgggtcccg aaagatacct tcgtcatggg gaagagtga tggagagaag 180
ataatataaa atgccaaacg tatataaatt gaaaggggaa aattgagaga acttgcgac 240
catggtcttg gtcttccata aagggggaact gggacatcca ctggagcaga gcacagattg 300
gcccaagagc cccaagactc cactggcct ccgcagaggc cgacagtgt tctgtcctgc 360
ggagattgtg gtttccctgt tagaaggaga ggagaacacc tgtggcaaac agaaaccaa 420
ggaaaacaat ttaaagccaa aatttcaggc ttctcaaggga gtaggtgtc tatatgaaa 480
ggagtcaatg aaaaaatcct tgaaagacag tgttgccctc acaataaag atcagaattc 540
catgaaacat gaggatccca gtatcatatc catggaagat ggggtcccat atgttaatgg 600
ctcattaggt gaagtgactc catgccaaca tgcaaagaag gcgaatggcc caaactatat 660
tcagcctcaa aaaagacaga ccacttttga aagccaggat cgcaaggcag tgtcccctag 720
cagttctgaa aagagaagta agaactctat ttctaggcca ttagaaggta agaagtcctt 780
aagtcttagt gcaaagactc acaacatagg ctttgacaaa gacagctgcc atagttacc 840
aaagacagaa gcttcacagg aagagcgggtc tgattcaagc ggctccatc ctctcaagaa 900

```

```

atcaccaaaag gtctcatcca aggcactcgg ggaaatcaaa actgatttct cactttctat 960
tagtaattcg tcagatgtga gtgctaaaga taagcatgct gaagacaatg agaagcgttt 1020
ggcagccttg gaagcaggc aaaaagcaaa agaagtgcag aagaagctgg tgcataatgc 1080
tctggcaaat ttggatggtc atccagagga taagccaacg cacatcatct tccgttctga 1140
cagtgaatgt gaaacagagg agacatcgac tcaggagcag agccatccag gagaggaatg 1200
ggtgaaagag tctatgggta aaacatcagg gaagctgttt gatagcagtg atgatgacga 1260
atctgattct gaagatgaca gtaatagggt caaaattaaa cctcagtttg agggcagagc 1320
tggacagaag ctcatggatt tacagtgcga ctttggcacc gatgacagat tccgcatgga 1380
ctctcgattt ctagaaactg acagtgaaga ggaacaggaa gaggtaaatg aaaagaaaac 1440
tgctgaggaa gaagagcttg ctgaagaaaa aaagaaaagcc ctgaatgttg taaaaagtgt 1500
tttgcaaatc aacttaagca attctacaaa cagaggatca gtagctgcta agaaatttaa 1560
ggacatcata cattatgatc caacgaagca agaccatgcc acttacgaaa gaaaaagaga 1620
tgataaacca aaagaaagta aagcaaaacg aaaaaagaaa agggagggaag ctgagaaact 1680
acctgaggtg tctaaagaaa tgtattataa tattgctatg gatctgaaag aaatattcca 1740
aactacaaaa tataccagtg aaaaggaaga gggcacaccc tggaaatgagg actgtggtaa 1800
agagaaacct gaggaaatcc aggaccctgc agctctgacc agtgacgctg agcagcccag 1860
cgggttcacg ttctcttttt ttgattcaga cactaaagac ataaaggaag agacctacag 1920
agttgaaaca gtgaaacctg gaaagattgt ctggcaggaa gacctctgtt tacaagacag 1980
cagttcagaa gaggaagatg ttactgaaga aacagatcac agaaactcca gtcctggaga 2040
agcatcatta cttgagaaag agaccactag atttttcttt ttctctaaga atgatgaacg 2100
acttcaaggt tctgacttat tctggagagg agtaggaagt aatatgagca ggaactcttg 2160
ggaggccaga acaaccaacc tgcgtatgga ttgtcgaaag aaacataaag acgcaaaaag 2220
gaaaatgaaa ccaaaataat aaatgtcagc tgggttttgat actgaatgtg aacaaggctc 2280
acctaaggaa actgaccacg aaaacagttt tagctgacaa agaagaaatt tcagagtga 2340
ggaattttta aaatctggct gacggaatat cattctgggt gccatctttt tctgtggaac 2400
tcctctgcag ttcttcttaa gtaattactt caaaaattaa attcaacttc ttataaagga 2460
agaacaagat agtccttgaa aatacttttt gtatataatc tctttgccct ctatcctgag 2520
taactaatgg acatcttctc atgcaagggt tatatgaagc ctttttaaat aaatgagtca 2580
aagcaaaaaa aaaaaaa 2597

```

<210> 39
 <211> 2641
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4586187CB1

```

<400> 39
ctgggaagaa agctatcagc accaactcag aactctccac cttcagatca gacattctag 60
atctccgtca gcaacttcgt gagattacag aaaaaaccag caagaacaag gatacgctgg 120
agaagttaca ggcgagcggg gatgctctgg tggacaggca gagtcaattg aaagaaactt 180
tggagaataa ctctttcttc atcaccactg taacaaaaac cctccaggcg tataatggct 240
atgtcacgaa tctgcagcaa gataccagcg tgcctcaggg caatctgcag aaccaaattg 300
attctcataa tgtggtctac atgaacctca acaacctgaa cctgaoccag gtgcagcaga 360
ggaacctcat cacgaatctg cagcggctct tggatgacac aagccaggct atccagcgaa 420
tcaagaacga ctttcaaaat ctgcagcagg tttttcttca agccaagaag gacacggatt 480
ggctgaagga gaaagtgcag agcttgacga cgctggctgc caacaactct gcgttggcca 540
aagccaacaa cgacacctg gaggatatga acagccagct caactcttc acaggtcaga 600
tggagaacat caccactatc tctcaagcca acgagcagaa cctgaaagac ctgcaggact 660
tacacaaaga tgcagagaat agaacagcca tcaagttcaa ccaactggag gaacgcttcc 720
agctctttga gacggatatt gtgaacatca ttagcaatat cagttacaca gccaccacc 780
tgcggaacgt gaccagcaat ctaaatagaag tcaggaccac ttgcacagat acccttacca 840
aacacacaga tgatctgacc tccttgaata ataccctgac caacatccgt ttggattctg 900
tttctctcag gatgcaacaa gatttgatga ggtcagggtt agacactgaa gtagccaact 960
tatcagtgat tatggaagaa atgaagctag tagactcaa gcatggctcag ctcatcaaga 1020
attttacaat actacaaggt ccaccgggcc ccagggttcc aagaggtgac agaggatccc 1080
agggaccccc tggcccaact ggcaacaagg gacagaaagg agagaagggg gagcctggac 1140
cacttgcccc tgcgggtgag agaggcccaa ttggaccagc tgggtcccccc ggagagcgtg 1200
ggcgcaaaag atctaaggc tcccagggcc ccaaaggctc ccgtgggttc cctggggaagc 1260
ccggccctca gggccccagt ggggacccag gccccccggg cccaccaggc aaagaggggac 1320
tccccggccc tcagggccct cctggcttcc agggacttca gggcaccgtt ggggagcctg 1380
gggtgcctgg acctcgggga ctgccagggt tgcctggggt accaggcatg ccaggcccca 1440
agggcccccc cggccctcct ggcccatcag ggcgggtggt gccctgggcc ctgcagaatg 1500
agccaacccc ggacccggag gacaatagct gcccgctca ctggaagaaac ttcacagaca 1560
aatgctacta tttttcagtt gagaaagaaa tttttgagga tgcaaaagctt ttctgtgaag 1620

```

```

acaagtccttc acatcttggtt ttcataaaca ctagagagga acagcaatgg ataaaaaac 1680
agatggtagg gagagagagc cactggatcg gcctcacaga ctcagagcgt gaaaatgaat 1740
ggaagtggct ggatgggaca tctccagact acaaaaattg gaaagctgga cagccggata 1800
actggggctca tggccatggg ccaggagaag actgtgctgg gttgatttat gctgggcagt 1860
ggaacgattt ccaatgtgaa gacgtcaata acttcatttg cgaaaaagac agggagacag 1920
tactgtcate tgcattataa cggactgtga tgggatcaca tgagcaaatt ttcagctctc 1980
aaaggcacaag gacactcctt tctaattgca tcaccttctc atcagattga aaaaaaaaaa 2040
gcactgaaaa ccaattactg aaaaaaaaaa gacagctagt gttttttacc atccgtcatt 2100
acccaaagac ttgggaacta aaatgttccc cagggtgata tgctgatttt cattgtgcac 2160
atggactgaa tcacatagat tctcctccgt cagtaaccgt gcgattatac aaattatgtc 2220
ttccaaagta tggaacactc caatcagaaa aaggttatca ttggctcgtt agttatggga 2280
agaactttaag catatactgt gtaaacagtg ccatacattt ctaaaatccc aagtgtagga 2340
aaaatatgca gacatacaga tatataggcc aactattagt aataatatga aatatactta 2400
aagagctttt aaaactttgt atttttgtac aaaatatttg tctttttaca tttttttcct 2460
tttttttttt ttgtcatttt accgacataa tacatggagc caaagaaaaa aataatggta 2520
ctaataaaaa ctcttagggg ttcctgtcag atttaattct acccagtggc aaagaatttt 2580
ttcaattgtg gctttaaaaa aataattaaa tatacatgta tatatatata aaaaaaaaaa 2640
a

```

<210> 40
 <211> 914
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 401801CB1

```

<400> 40
cagaggtgta ctctacatga tgcacaaatg tgatatgtct ttttggtgaa gagcacgtga 60
tgaaaacagt aaataactga gaatagcaca aagctactag ggactcagat gattcaaata 120
ttgaagacta tgagtaatat taatcagcaa cttagtttgt tatcttcagt tatatgggag 180
gagtacatta ttctttgtta taggactacc tcacccttaa attgtaagtt ctttattagc 240
ttattgcatt ttccattctt aaagcagtag tttagtgttc tcttactgta tgacaattaa 300
aagtatactt aattgactac ataatgtgat agttaaaaaa tatatttaaa gtagcttttt 360
gaaagctttg ctgtttcccc ccctttttgt atataaaagc attagtgttc attattcatg 420
tgttctcatt actttttaca aatgagaaca atgccttatg catgttgtga acaattacta 480
aaatttttatg taaatttaac ctttattttt aattaatatg tgtaagata taacattatt 540
ttatctatta atatatgtat ttaattttac agggaaaacat ctttatgtat ggaggcagaa 600
ttgaaacaaa tgatggcaat gtcacagatg aattatgggt ttttaacata catagtcagt 660
catggagtac aaaaactcct actgttcttg gacatggtca gcagtatgct gtggagggac 720
attcagcaca tattatggag ttggatagta gagatgttgt catgatcata atatttggat 780
attctgcaat atatggttat acaagcagca tacaggaata ccatatctgt gagttactta 840
aaaattgtaa tttctttatt gattgggaat gtttttctct ttaataaaat cttcatatga 900
atttaaaaaa aaaa
914

```

<210> 41
 <211> 1006
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1721842CB1

```

<400> 41
ggcaaagaga actacaaatc ccagcgtcac ccgcggcctt gaagccccgc cctgacaaa 60
ctgaaggtcc cggtaagcat cgcgtcagta cttatggcgc ctgccgggtt gtggtgacga 120
aagcagttgc catggagttg ctctgagtaa ccttgaggca gtgggacgcc aagactggag 180
aggaagcgac tgcggggagt atttccattt taaccggaaa caatccctga acccagga 240
atgaatgcct aatggtggag ttccagccat cagtgcacag ctgaaccag actcccagg 300
cacctgcttg cacctttgaa tgatggcctg aactatgaac aaacgggact atatgaacac 360
ttcggtagag gagccccctc ttgactactc cttcagaagc atccacgtca ttcaagatct 420
ggtaaatgag gagccaagga caggactacg accactgaag cgttcaaagt cggggaaatc 480
actgacccag tccctgtggc tgaataacaa tgttctcaat gatctgagag acttcaacca 540
ggtggcttca cagctgttgg agcaccacga gaacctggcc tggatcgacc tgtcctttta 600
tgacctgact tccattgacc ctgtcctaac aactttcttc aacctgagtg tctctatct 660

```

tcacggcaac	agcatccagc	gcctggggga	gggtgaataag	ctggctgtcc	ttcctcggct	720
ccgtagcctg	acactccatg	ggaaccccat	ggaggaagag	aaaggggtata	ggcaatatgt	780
gctgtgcacc	ctgtcccgtg	tcaccacgtt	cgacttcagt	ggggtcacca	aagcagaccg	840
caccacagct	gaagtctgga	aacgcatgaa	catcaagccc	aagaaggcct	ggaccaagca	900
gaatacactt	tgaggctccc	acgaccctag	tagtcctaaa	ggcctaagca	tagacagcat	960
ggtttgacaa	taaataattt	gagctgttga	gcagaaaaaa	aaaaaa		1006

<210> 42

<211> 2582

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1833221CB1

<400> 42

gttaaat	gtactgaatg	cagccttttg	ttgtttaaaa	aatttttttg	aactcacagt	60
ctacatcagc	atcagcatct	gcgtcaccat	ttcaatctgc	atgggtatagt	gaatctgaga	120
taactcaggg	agcacgctca	agatcgcaga	accagcaacg	ggatcatgat	tcaaaaagac	180
ctaaactttc	ctgtacaaac	tgtactacct	cagctgggag	aaatggttga	aatggtttaa	240
acacattatc	agattcatct	tggaggcata	gtcaagttcc	tagatcttca	tcaatgggtac	300
ttggatcatt	tggacacagac	ttaatgagag	agaggagaga	tttggagaga	agaacagatt	360
cctctattag	taatcttatg	gatttatagtc	accgaagtgg	tgatttcaca	acttcatcat	420
atgttcaaga	cagagttcct	tcataattcac	aaggagcaag	acaaaaagaa	aactcaatga	480
gcactttaca	gttgaatata	tcattccacaa	accaccaatt	gccttctgaa	catcagacca	540
tactaagttc	tagggactcc	agaaaattcct	taagatcaaa	tttttcttca	agagaatcag	600
aatcttcccg	aagcaatacg	cagcctggat	tttcttacag	ttcaagtaga	gatgaagccc	660
caatcataag	caattcagaa	aggggtgttt	catctcaaaag	accatttcaa	gaatcttctg	720
acaatgaagg	taggcggaca	acgaggagat	tgctgtcacg	catagcttct	agcatgtcat	780
ctactttttt	ttcacgaaga	tctagtcagg	attccttgaa	tacaagatca	ttgaattctg	840
aaaattctta	cgtttctcca	agaatcttga	cagcttcaca	gtcccgtagt	aatgtaccat	900
cagcttctga	agttcccgat	aataggggcat	ctgaagcttc	tcagggattt	cgatttctta	960
ggcgaagatg	gggtttgtca	tctcttagcc	acaatcatag	ctctgagtc	gattcagaaa	1020
attttaacca	agaatctgaa	ggtagaaaat	caggaccatg	gttatcttcc	tcacttagaa	1080
atagatgcac	acctttgttc	tctagaagga	ggcgagaggg	aagagatgaa	tcttcaagga	1140
tacctacctc	tgatacatca	tctagatctc	atatttttag	aagagaatca	aatgaagtgg	1200
ttcaccttga	agcacagaat	gacccctctg	gagctgctgc	caacagacca	caagcatctg	1260
cagcatcaag	cagtgccaca	acaggtggct	ctacatcaga	ttcgggtcaa	ggtggaagaa	1320
atacaggaat	atcagggatt	cttccctggt	cttattccg	gtttgcagtc	ccccagcac	1380
ttgggagtaa	tttgaccgac	aatgtcatga	tcacagtaga	tattattcct	tcaggttggg	1440
attcagctga	tggtaaaagt	gataaaacta	aaagtgcgcc	ttcaagagat	ccagaaagat	1500
tgcagaaaat	aaaagagagc	ctccttttag	aggactcaga	agaagaagaa	ggtgacttat	1560
gtagaatttg	tcaaatggca	gctgcacat	catctaattt	gctgatagag	ccatgcaagt	1620
gcacaggaag	tttgcagtat	gtccaccaag	actgtatgaa	aaagtgggta	caggccaaaa	1680
ttaaactctg	ttcttcatta	gaagctgtaa	ccacctgtga	actatgtaaa	gagaagttag	1740
agcttaacct	ggaggatttt	gataattcat	aactacatag	agctcatgca	aatgaacaag	1800
ctgagtatga	gtttatcagc	tctgggtctc	acctagtggg	gttattgcac	ttgtgcgaac	1860
aaagcttttc	tgatatgatg	ggaaatacaa	atgaaccaag	cacacgtgtc	cgatttatta	1920
accttgcaag	aactcttcag	gcacatatgg	aagatctcga	aacttcagag	gatgattccg	1980
aagaagacgg	agaccataac	aggacatttg	atattgccta	acttcatata	agacagatgg	2040
atgatctgtg	aacataagtg	tttattaaaa	atggcaatta	aatataaatt	acttttgttg	2100
gggaatgcct	aataaataca	ttgactatat	ataaaatgaa	tatatacata	cacatgtatg	2160
cctgtatata	tatattcatt	ctccagtgtt	gctgaattaa	aattctgctg	gactttttta	2220
catagcaaat	ccgatgttta	taaactggta	atcaaaaagg	tttttctttt	taggtgagtg	2280
ggaaagtatt	acccttggtt	taaatatcta	agcaatgcct	atcaaccctt	ttttgtgtta	2340
tgattactgt	agtcataatt	atgaaaaaag	gtttgtgttt	tactcttgct	agtgagaaaa	2400
gtgggacaaa	atatactttt	gaaataaaat	gctatatggc	acctaattat	tttttctttt	2460
aaaatgcctt	aagttgcagt	ctcatttttg	taatcatttg	cttccagtgt	ttaaaaaatta	2520
aaaaaagaat	ggggagaagg	ttatgagaag	agcattatta	agtttccaaa	tttaatttga	2580
at						2582

<210> 43

<211> 2849

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2041168CB1

<400> 43

```
acgtgaccaa gagctgacgt gtgcagaagt ccttcttctg ctggctcgttg tccccgtctg 60
agtaccagct cccactgcc ctgagggcgg gccggcctgc ggcggaggga aaaaggaaga 120
ggagaaggaa attgtcccga atccctgcag tctttctgta ggttgccgga caacgccagg 180
caaaagaaga ggaaggaatt taatcctaata cgggtggaggt cgatttgagg gtctgtctgta 240
gcaggtgggt ccgcttgaag cgaggggagga agtttccctcc gatcagtaga gattggaaaag 300
attgttggga gtggcacacc actagggaata agaagaaggg gcgaactgct tgtcttgagg 360
aggccaaccc ccagaatcag ctcttgtggc cttgaagtgg ctgaagacga tcacctcca 420
caggcttgag cccagtcacca cagccttccct cccccagcct gagtgactac tctattcctt 480
ggctccctgct attgtcgggg acgattgcat gggctacgcc aggaaagtag gctgggtgac 540
cgcagcctgg tgattggggc tggcgcctgc tattgcattt atagactgac taggggaaga 600
aaacagaaca aggaataaat ggctgagggt ggatctgggg atgtggatga tgcctgggac 660
tggtctgggg ccagggtataa tgactggtct gatgatgatg atgacagcaa tgagagcaag 720
agtatagtat ggtacccacc ttgggctcgg attgggactg aagctggaac cagagctagg 780
gccagggcaa gggccagggc taccggggca cgtcgggctg tccagaaacg ggcttcccc 840
aattcagatg ataccgtttt gtccctccta gagctacaaa aggttctttt ctgtggttag 900
atgtctgaaa agccttatat tcttgaagca gctttaaattg ctctgggtaa caatgctgct 960
tatgcattta acagagatat tttcgtgat ctgggtgggt tcccaattgt cgaaagatt 1020
ctcaatactc gggatcccat agttaaggaa aaggctttaa ttgtcctgaa taacttgagt 1080
gtgaatgctg aaaatcagcg caggcttaaa gtatacatga atcaagtgtg tgatgacaca 1140
atcacttctc gcttgaactc atctgtgcag ctgtctggac tgagattgct tacaatatg 1200
actgttacta atgagtatca gcacatgctt gctaattcca tttctgactt tttctgctta 1260
ttttcagcgg gaaatgaaga aaccaaactt caggttctga aactcctttt gaatttggct 1320
gaaaatccag ccatgactag ggaactgctc agggcccaag taccatcttc actgggctcc 1380
ctctttaata agaaggagaa caaagaagtt attcttaaac ttctgggtcat atttgagaac 1440
ataaatgata atttcaaatg ggaagaaaaa gaacctactc agaatcaatt cgggtgaagg 1500
tcactttttt tctttttaaa agaatttcaa gtgtgtgctg ataaggttct gggaatagaa 1560
agtcaccatg attttttggg gaaagtaaaa gttggaaaat tcatggccaa acttgctgaa 1620
catatgttcc caaagagcca ggaataacac cttgattttg taatttagaa gcaacacaca 1680
ttgtaaacta ttcattttct ccacctgtt tatatggtaa aggaatcctt tcagctgcca 1740
gttttgaata atgaatatca tattgtatca tcaatgctga tatttaactg agttggctct 1800
taggtttaag atggataaat gaatatcact acttgctctg aaaacatggt tgttgctttt 1860
tatctcgtg cctagattga aatattttgc tatttcttct gcataagtga cagtgaacca 1920
attcatcatg agtaagctcc cttctgtcat tttcattgat ttaattttgt tatcatcaat 1980
aaaattgtat gttaatgctg gaaagaaaaa aagaagaaag aaaaaaacca tccctgtcct 2040
tcagtttata atctagttgg agagataaga aacgtacaaa ccaaaagata acagaatatc 2100
tgaagcatgt actcattgtc agatgttccc tctgagagca cagaggaggc aaaagcttct 2160
gtgggatgtg ctatgcggct aaagcttcac agaggaggtg gcaattgaaa atgagtcctg 2220
aatggggtag ggtggttagg gaattccatg agacaagaca aggggggcat ggtgtgagaa 2280
aggcatggaa gtaggaaccc tcttccatg acaggagatc attctgctta gaggggagag 2340
tgtggagagt gggagttagt aattttggaa agctgggtga agccagttgt ggagaattgt 2400
ttgaatatta tcccattgaa taccagagc cactaaatct ttttttacta gaaaataatt 2460
ggggtccata tgaaagtctc tattactgag tagtgtcaat gaggggtgtg caaaatggag 2520
cctttcacat cctagtgtg gccatttggt aatacagata taagccttaa actatgtaaa 2580
ccccgtcct aaggaagtaa ttgaataatt gcccaaagat tgtatgtatg aggctgttca 2640
tccagcact gtctaagcta gtaaaaattg gaaacaattt aagtatctag cacattggat 2700
tggttataaa gcaaggaatg ttcacacagt aggatattat aagtatgctg atggaaatct 2760
atattgccag gaaaagctat tcattatgct ttgtgaagtc agaaagtaaa aaagggtaga 2820
tagaagtatt cgaagtatag ttccatttt 2849
```

<210> 44

<211> 670

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2365794CB1

<400> 44

```
gaggcaagaa ttcggcacga ggggcgcgcg gggcatttct tccactgcc gtctgaggga 60
acgctaagta gtgtgtccgg cgcctgttcc cagctccgcg ttgttccgcg agaaagcgag 120
aggccgagcc cgggctgggt cgatggccgc ggtggtggcc aagcgggaag ggccgccgtt 180
```

```

catcagcgag gcgggccgtgc ggggcaacgc cggcgtcctg gattattgcc ggacctcggt 240
gtcagcgctg tcggggggcca cggccggcat cctcggcctc accggcctct acggcttcat 300
cttctacctg ctgcctcccg tctgtctctc cctgtccttc attctcaagg cgggaaggag 360
gtggaacaaa tatttcaaat cacggagacc tctctttaca ggaggcctca tcgggggcct 420
cttcacctac gtccgtgttct ggacgttcct ctacggcatg gtgcacgtct actgaaatgg 480
gggcccgggg gactttttta aaaaaccaga tcggggaggac tgtggccagc aattaacacc 540
atgtagactt ccttagttct taagtgggtg aattcgctgc ttgttctgta acgttataaa 600
taatttatat ctgaagacgg agagcctgta atattcttca gattaaatga agcgtgagac 660
aaaaaaaaaa                                     670

```

<210> 45
 <211> 2364
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2618452CB1

```

<400> 45
ctcaatgccacacaggcacca ttcctccacc gacaacgctg aaggccacag ggtccacca 60
cacagcccca ccaatgatgc caaccaccag tgggaccagc caagcctcaa gtccttcaa 120
cacagccaaa acctctacat ccctacattc acacacttcc tccacacacc atcctgaagt 180
caccccaact tctatcacca acatcacccct caacccccacc agtataggaa cctggacacc 240
cgtggcccac accacctcgg ccaccagcag caggctaacc acacccttca ccacacattc 300
cccactaca gggagcagtc ccatctcttc cacaggctct atgactgcaa catcctttca 360
gaccaccact tattatacac caccatcaca cctcagacc acacttccca ctacgttcc 420
acctttctcc acctccttgg tgactccaag tactcacaca gtcacatca ctaccacac 480
acagatggcc acttctgcct ccattccactc aacgccaaaca ggcaccgttc ctccaccaac 540
aacgctcaag gccacagggt ccacccacac agccccacca atgacagtga ccaccagtgg 600
gaccagccaa acccacagct cattcagcac agctacagcc tcttcttct tcatatectc 660
ctcgtcttgg tcgtcttggc tgccctcagaa ctctagctca aggccaccgt catcacctat 720
caccacacaa ctccccact tgagttctgc aaccactcct gtttccacaa ctaatcagct 780
gtcctcctca ttttctccca gtccttctgc cccctctact gtttcttctt atgtgcctc 840
ctccccactc tctccccaga cttcatcgcc ttctgttggc acatcttctt ctttctgtgc 900
cgcccccggt cactccacaa ccctgagctc ggggtcacac tctcattgt ccactcatcc 960
cacgactgca tcagtgtctg catctcctct ttttcttct tctccagctg cctctactac 1020
cattagggcc actctcccc acactatctc ctctccttcc accctctctg ctctactccc 1080
catatccact gttaccgtgt ctcccacccc atccagccac ctagcctcca gcaccattgc 1140
atttccgctc attcccgagg ccacggccag caccacacac gccctgcct tctctctca 1200
gtccaccacc tcgcggtcca cttctctcac caccgagtt cccacatcag gctttgtgtc 1260
actcacctcg ggggtgacgg gtatccccac ctctccagtc accaaccctta ccaccaggca 1320
ccctgggtccc acctgtgcgc ctaccacacg gtctctgacc agctccctca ctgccatgg 1380
aagcacccct gcttctgccc cggtatcttc tctcgggaca cctacgccc cctcaccgg 1440
ggctctgagc gtgcgggagc agcaggagga gatcacgttc aaggggtgca tggcgaacgt 1500
gacggtaacc gcctgtgagg gcgcctgcat ttccgctgcc agcttcaaca tcatacca 1560
gcagggtgat gcccgctgca gctgtgcgcg cccctccac tcctatgagc agcagctgga 1620
gctgccttgc cccgatccca gcacgcctgg ccggcggtc gtactcacc tgcaggtgtt 1680
cagccactgc gtgtgcagct ctgtggcctg tgagactag cagggtcgt gctgtctctc 1740
ctggggctga aggactgcag atgacagaca ggaaaacacc caccagccc cttccgctt 1800
gtgccagcag ctgctttctt ggtcaccagg cctggcccc aagtgcctg ggcgtgggt 1860
ccctggggca ccggttggag aggggtgcc aagcaggggc tcagactacc acactcctgc 1920
agaccctgag ccagcagaga gggactgagg cggacagtgg tcacggacct ccaggcaca 1980
cagggcactc ccgaccaccc ctgcccactg tccaacacct ccagccctt gaacttggcc 2040
ccagccctgc tgggcccaga accctgcaga tgaagccaca gagcaggcgc tcgaccagac 2100
ccatcagggg cgaggagggc acggaaacct gtgccgagat gggggcaaga ggcccaggca 2160
gccaccagca cagagaagag gagatcccca gagtcaggga gggcagaggg tggcagcgag 2220
ggcagggcag ccgccccgcg tcccagccag gcagaaggcc cccaccagca ccacacccat 2280
ccccagcagc ctgtccttgg gagagggcgt cacccggtca gagactccaa ataaaccggt 2340
tcttgtcaag gcacaaaaaa aaaa                                     2364

```

<210> 46
 <211> 3600
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2622288CB1

<400> 46

gcagaggcgg	cgggggtcct	cctcccgcgc	ctcctcggcc	tccccctcgg	gcgctctcgc	60
gctaactgtg	ctcctccggg	gccctccgcc	tgctcccagc	catgggtggc	tgggcgctcgg	120
cggttcctgt	ctgcctcgct	ttctccttgg	ccaccctggg	ccagcgagga	tctgggggact	180
ttgatgattt	taacctggag	gatgcagtga	aagaaacttc	ctcagtaaag	cagccatggg	240
accacaccac	caccaccaca	accaataggc	caggaaccac	cagagctccg	gcaaaacctc	300
caggtagtgg	attggacttg	gctgatgctt	tggatgatca	agatgatggc	cgcaggaaac	360
cgggtatagg	aggaagagag	agatggaacc	atgtaaccac	cacgaccaag	aggccagtaa	420
ccaccagagc	tccagcaaat	actttaggaa	atgattttga	cttgggtgat	gccctgggatg	480
atcgaaatga	tcgagatgat	ggccgcagga	aaccaattgc	tggaggagga	ggttttttcag	540
acaaggatct	tgaagacata	gtaggggggtg	gagaatacaa	acctgacaag	ggtaaagggtg	600
atggccggta	cggcagcaat	gacgaccctg	gatctggcat	ggtggcgagag	cctggcacca	660
ttgccggggg	ggccagcgcc	ctggccatgg	ccctcatcgg	tgccgtctcc	agctacatct	720
cctaccagca	gaagaagttc	tgcttcagca	ttcagcaggg	tctcaacgca	gactacgtga	780
agggagagaa	cctggaagcc	gtggtatgtg	aggaacccca	agtgaataac	tccacgttgc	840
acacgcagtc	tgacagagccg	cgcgcgcgcg	ccgaaccagc	ccggatctga	gggcccctgtc	900
cagctgcagg	catgcacaat	ggtgccaccg	cttgctaccc	ggctcccccc	accccttcat	960
ttggacccgc	agctgctgtg	ctgctctgtg	ccatcggtct	cttggttggtc	tgagtttccc	1020
ggatgagctc	tgggtgtttg	tgagtttggg	ttctctgccc	tgccccaaagc	gtgctgagac	1080
ttggtgccga	aattcaagag	ccagctctga	tagaaagcca	gcaccagcct	cgggagctgc	1140
tgagccacca	actcccaaag	ccagcctgcc	tccagcttta	ctgagcacag	gatgcggggg	1200
ccaagatgat	gctgaggcct	gatgacattt	atgcttaggg	gacaagagtt	tgaactcaag	1260
ggactgtgac	ccctgcacac	tggagtggct	catgtgtgca	ggtttctgcc	aatagacagc	1320
ccctgacagt	ggcctcaagg	agctgcaggt	gggggggtca	gcttgcaccc	acttggagcc	1380
cctgcaagga	gcgaaccggg	cagcaccaag	taacaccaca	cacacgcagc	accaggatg	1440
atggtttcac	ttcagttctc	cccattcccag	gtttttatgtt	gctgggcttc	cggagagccg	1500
gtccaagcgc	aggctttcag	tgatttaagt	acaacaatgc	atctcgtgat	agtcctgcct	1560
tgagagctta	ggaaatcttc	ggataagtat	gaagcaattc	gtaggcctgt	ttcccattctg	1620
attccatagg	gggctgggtg	tggccttcgg	gttgacatga	gaaaggctctt	tagcaatcat	1680
ttctgcaccg	gagatgagtt	ttatcctgtg	ttgggggagag	gtgctcacc	tccaccctgt	1740
gtccctgttt	tggtagcaag	agtgaccgat	gtcaagaacg	agcatcaaa	ccagaatcct	1800
gcttgttttg	ttaaaaatgt	aattgggggc	ggcgggggag	gagaggggaa	agagacattc	1860
gcttgggtta	gtgaaacgca	ggtgactttg	tagctctgtg	gtcagcctac	ttgtctgctc	1920
tgagggagag	tgctgtggga	gccatgctca	ccgtggcaaa	cacaggaacc	ccatgactcg	1980
ccccacacct	ggcgtggagc	tgccctgggtt	gggctgggagc	agagctgggt	tccctggaatg	2040
ttcctttggc	ccacatatgg	ttctgtcccc	gtgagctctg	ttgtcagagg	ctcacgggac	2100
agaaccacat	gctaggggtc	agggccctg	ttactgata	gtcagtttgc	tgtgtcagaa	2160
agcacttctg	aaagcagata	tgagtcacca	gacaggcagg	atcttacaaa	actcacgggc	2220
ctcttttggt	tgcatgatgg	ccccatgcgt	ttcataggct	gtccactgag	cgggattgtc	2280
tgctgagtg	gatgagccaa	ctccagtttc	ttaaggaaac	caactggaatc	tgacgcccc	2340
acatgcatct	gtctaacgca	tgccctgtgt	tcgttttgca	aacatgcctg	tgggtggagg	2400
tggtcagttg	tagccctgtg	cgctctaggg	ctgctctgtg	aggccattcc	cagtgcgtgc	2460
ccttgagctc	cttaccaccc	cttttctctg	tcggcccttt	aatccctgac	agacctggac	2520
tgtgtggctg	aagggggacc	tgacgactg	cagaaatgcc	tctgcgtggg	gccatgaagg	2580
aaagaaacct	tggcctgggtc	tcgagaagct	tcccatgctt	cagggaagta	gtaagggtgg	2640
ggtggcttgc	aggattggcc	tgtttccagg	gctcccaca	ctcattggcc	agattgtgaa	2700
ctttgtcagg	cttgccctc	cctgatacca	agtatgtcga	gaaccgatgg	ccccaccctc	2760
tggctgggtg	tgggcccggg	gtggctatgg	aggattttgg	catgcgtggc	ctgtcgccac	2820
ctggacagcg	tgacctcagg	ggttgtccac	tttaccttta	tgggtgaggcc	tgctcgatgg	2880
ctaagtcctt	gaaaccctag	agctgtgacg	tagaatatgt	gctgtctgtg	agacctgtgt	2940
cccaggagca	ctgactgcag	ttgagagaga	cccattttgc	tctcccttac	cgccccccgc	3000
cccgggtgct	ttctgcacaa	agcctagagc	ctggcactca	agcccaccgg	tggcagctcc	3060
tagtgactgg	acatgcctgg	aagacccttc	agccttctgt	ttgcagaacg	ttcattttcag	3120
gagcttctcc	ttcccacaga	catcttacac	ttgctcgaca	ctgccacctg	cagaagcctg	3180
gcgggctctg	gtcaccatgt	gtctatctga	aggttgactt	ggccagcatg	ggcctgtccc	3240
aagcgagagg	ggagcacacg	tggactgaaa	ggactgggtg	aaagtggcca	atctctatca	3300
gcttaatttg	gcagagaaaa	tttgtaacaa	ctctgagcac	atgctgggtg	aagtccacagc	3360
tcaaggaaag	ataaagctgg	gcggaaggag	gtgtgcgtgg	cttctggggg	gggaccacaga	3420
ggggaggctc	tgggacaggg	gctgggggtc	agtgccaggg	ccctgaggaa	gaaatgggga	3480
ctgatctcaa	aattccagaa	ttccctgtac	atctgttcac	gtgcttgtgt	ccaggtgtga	3540
cttgtaaact	gtctagtgtt	tgcattaaat	aaaatggcac	cgagcagaaa	aaaaaaaaaa	3600

<210> 47

<211> 1236

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2806595CB1

<400> 47

ttaatttccc	cgaaatcaga	ctgctgcctt	ggaccgggac	agctcgcggc	ccccgagagc	60
tctagccgtc	gaggagctgc	ctggggacgt	ttgccctggg	gccccagcct	ggccccgggc	120
accctggcat	gaggagatgg	gcctgttgct	cctgggtccg	ttgctcctgc	tgccccggctc	180
ctacggactg	cccttctaca	acggcttcta	ctaactcaac	agcgccaacg	accagaacct	240
aggcaacggt	catggcaaag	acctccttaa	tggagtgaag	ctgggtgggtg	agacacccga	300
ggagaccctg	ttcacctacc	aagggggccag	tgtgatcctg	ccctgccgct	accgctacga	360
gccggccctg	gtctccccgc	ggcgtgtgcg	tgtcaaatgg	tggaagctgt	cggagaacgg	420
ggccccagag	aaggacgtgc	tgggtggccat	cgggctgagg	caccgctcct	ttgggggacta	480
ccaaggccgc	gtgcacctgc	ggcaggacaa	agagcatgac	gtctcgtctg	agatccagga	540
tctgcggctg	gaggactatg	ggcgttaccg	ctgtgagggtc	attgacgggc	tggaggatga	600
aagcgggtctg	gtggagctgg	agctgcgggg	tgagatgcta	acggggactg	ggtgacactg	660
ggacctgaga	gcagagggga	gaggaccaga	gaaaacatcc	agacctctgt	gcttttagaca	720
tttaaaagta	cttaattctc	aaaacaaccc	acatggaagc	tactgtttgt	acccccattt	780
tacaggtgag	aaaactgagg	cacagagagg	tcaagtaact	tacctaaggt	cacacagctt	840
gtaaaccgaca	gagctggggg	ctgaacccaa	gcacccagcc	tctagaatct	gttccccctc	900
accgcgtgta	attcacatct	cattcagaga	gaggaaaacc	agagctgggtc	ccacagctta	960
ttagagacag	agctgagatt	taagcaaggt	tagcgggttaa	cacaagcgaa	tgaggcagcc	1020
cactgtgaga	cgatgggagt	gggtgggagt	gggcacactc	ctgagccctt	gtcctgtgcc	1080
cagggagctc	ccacactact	caggcccagc	tgatgatgcc	acgccagaat	gcagacccca	1140
ctgccaaatt	ttctggcttg	acaagataag	cttatatttt	tatgtgagat	ctccagattt	1200
ttatgtaaaa	acctcctttt	taaaaacaaa	acaaaa			1236

<210> 48

<211> 3081

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2850987CB1

<400> 48

gccccggcca	cgccggcgcc	ggccggcgcc	gagagagctg	gctcagggcg	tccgctagcc	60
tcggacgacc	tgctgagcct	cccaaaccgc	ttccataagg	ctttgccttt	ccaacttcag	120
ctacagtgtt	agctaagttt	ggaaagaagg	aaaaaagaaa	atccctgggc	cccttttctt	180
ttgttctttg	ccaaagtctg	cgttgtagtc	tttttgccca	aggtctgtgt	gttttttagag	240
gtgtctatctc	cagttccttg	cactcctgtt	aacaagcacc	tcagcgagag	cagcagcagc	300
gatagcagcc	gcagaagagc	cagcgggggtc	gcctagtgtc	atgaccaggg	cgggagatca	360
caaccgccag	agaggatgct	gtggatcctt	ggccgactac	ctgacctctg	caaaattcct	420
tctctacctt	ggtcattctc	tctctacttg	gggagatcgg	atgtggcact	ttgcgggtgtc	480
tgtgtttctg	gtagagctct	atggaaacag	cctccttttg	acagcagttc	acgggctggg	540
gggtggcagg	tctgttctgg	tctggggagc	catcatcggg	gactgggtgg	acaagaatgc	600
tagacttaaa	gtggcccaga	cctcgtctgg	ggtacagaat	gtttcagtca	tctgtgtgtg	660
aatcatcctg	atgatgggtt	tcttacataa	acatgagctt	ctgacctgt	accatggatg	720
ggttctcact	tctgtctata	tctgtatcat	cactattgca	aatattgcaa	atttggccag	780
tactgctact	gcaatcacia	tccaaaggga	ttggattgtt	gttgttgtag	gagaagacag	840
aagcaaaacta	gcaaatatga	atgccacaat	acgaaggatt	gaccagttaa	ccaacatctt	900
agcccccatg	gctgttggtc	agattatgac	atgttggtcc	ccagtcatcg	gctgtggctt	960
tatttcggga	tggaaacttg	tatccatgtg	cgtggagtac	gtcctgtctc	ggaagggtta	1020
ccagaaaacc	ccagctctag	ctgtgaaaagc	tgggtcttaa	gaagaggaaa	ctgaattgaa	1080
acagctgaat	ttacacaaag	atactgagcc	aaaaccctcg	gagggaaactc	atctaattggg	1140
tgtgaaagac	tctaacaatc	atgagcttga	acatgagcaa	gagcctactt	gtgcctccca	1200
gatggctgag	cccttccgta	ccttccgaga	tggatgggtc	tctactaca	accagcctgt	1260
gtttctgggt	ggcatgggtc	ttgctttcct	ttatatgact	gtcctgggct	ttgactgcat	1320
caccacaggg	tacgcctaca	ctcaggggact	gagtgggtcc	atcctcagta	ttttgatggg	1380
agcatcagct	ataactggaa	taatgggaac	tgtagctttt	acttgggtac	gtcgaaaatg	1440
tggtttgggt	cggacaggtc	tgatctcagc	attggcacag	cttctctgtt	tgatcttgtg	1500
tgtgatctct	gtattcatgc	ctggaagccc	cctggacttg	tccgtttctc	cttttgaaga	1560
tatccgatca	aggttcattc	aaggagagtc	aattacacct	accaagatac	ctgaaattac	1620

aactgaaata	tacatgtcta	atgggtctaa	ttctgcta	attgtcccgg	agacaagtcc	1680
tgaatctgtg	cccataatct	ctgtcagtc	gctgtttgca	ggcgtcattg	ctgctagaat	1740
cggctcttgg	tcctttgatt	taactgtgac	acagttgctg	caagaaaatg	taattgaatc	1800
tgaagagggc	attataaatg	gtgtacagaa	ctccatgaac	tatcttcttg	atcttctgca	1860
tttcatcatg	gtcatcctgg	ctccaaatcc	tgaagctttt	ggcttgctcg	tattgatttc	1920
agtctccttt	gtggcaatgg	gccacattat	gtatttccga	tttgcccaaa	atactctggg	1980
aaacaagctc	tttgcttgcg	gtcctgatgc	aaaagaagtt	aggaaggaaa	atcaagcaaa	2040
tacatctgtt	gtttgagaca	gtttaactgt	tgctatcctg	ttactagatt	atatagagca	2100
catgtgctta	ttttgtactg	cagaattcca	ataaatggct	gggtgttttg	ctctgttttt	2160
accacagctg	tgcccttgaga	actaaaagct	gtttaggaaa	cctaagtcag	cagaaattaa	2220
ctgattaatt	tcccttatgt	tgaggcatgg	aaaaaaaaat	ggaaaagaaa	aactcagttt	2280
aaatacggag	actataatga	taacactgaa	ttcccttatt	tctcatgagt	agatacaatc	2340
ttacgtaaaa	gagtgggttag	tcacgtgaat	tcagttatca	tttgacagat	tcttatctgt	2400
actagaattc	agatatgtca	gttttctgca	aaactcactc	ttgttcaaga	ctagctaatt	2460
tatttttttg	catcttagtt	atttttaaaa	acaaattctt	caagtatgaa	gactaaattt	2520
tgataactaa	tattatcctt	attgatccta	ttgatcttaa	ggtatttaca	tgatagtgga	2580
aaaacaaaac	aatttaactag	aattctctaa	taaggtttat	ggtttagctt	aaagagcacc	2640
tttgattttt	tattatcaga	tggggcaaca	tattgtatga	agcatatgta	gcacttcaca	2700
gcatgggtat	catgtaagct	gcaggtagaa	gcaaagctgt	aaagtagatt	tatcacacaa	2760
tgactgcata	cagacttcaa	atatgtcaat	agtttggtca	tagaacctag	aagccaaaag	2820
ccacacagaa	gggcaagaat	cccaatttaa	ctcatgttat	catcattagt	gactctgtgt	2880
ctagaacatg	agggtgtaag	ccttcagcct	ggcaagttac	atgtagaaag	cccacacttg	2940
tgaagggttt	gttttacaaa	tcacttgatt	taacacactc	aggtagaata	tttttatttt	3000
tactgtttta	taccagaaag	ttatttctac	attgttctac	agcaagaata	ttcataaagg	3060
tggaccttgc	aagtgcgtat	a				3081

<210> 49

<211> 1825

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3557211CB1

<400> 49

cgtgttgaag	gcatacagacc	ctgacactga	ggacgatcag	ataatcttta	aaattcttaca	60
aggcccaaaa	catggacatc	tggagaacac	aacaacaggt	gaatttatcc	atgagaaatt	120
tagccaaaag	gacttaaaaca	gtaagactat	tctttacatc	ataaaacctat	ctttggaagt	180
aaattcagat	accgtggaat	ttcaaatcat	ggacccca	gggaactcgg	ccactcctca	240
aatttttgaa	ctgaagtggg	ctcatattga	atggtcacag	accgaatata	tctgtgagaa	300
tgtgggtttg	ttgcccttgg	aaattatcag	aaggggatat	tccatggact	cggcctttgt	360
gggtataaag	gtcaaccaag	tgtcagctgc	agttggaaaa	gatttcaccg	tgattccatc	420
taaaactgatt	cagtttgacc	caggaatgtc	aactaagatg	tggaatatag	caattaccta	480
tgacggatta	gaggaagatg	atgaggtcct	tgaagtaatt	ctgaactccc	ctgtgaatgc	540
agttcttggc	acaaagacaa	aagctgcagt	gaaaattttg	gactcaaaaag	gaggacaatg	600
ccatccttca	tattcctcca	accaaagcaa	gcacagcaca	tgggagaagg	gcatttggca	660
tctgtgccc	ccagggtcct	cctcatccac	cacttctggt	tcctttcatc	tggaaagaag	720
acctcttcca	tcttccatgc	agctagcagt	catcagggga	gacaccctgc	ggggctttga	780
tctacagat	ctttctcaaa	ggaagcttag	gacccgtggg	aatggcaaaa	cagttcgtcc	840
atcctctgtt	tatagaaatg	gaacagacat	catctataat	tatcatggga	tagtttcctt	900
gaaactggag	gatgacagtt	tcccaactca	caaaaggaag	gccaaagtat	ccatcattag	960
tcagccacaa	aagacaatca	aagtggcaga	actgcctcaa	gcagataagg	tggaaatccac	1020
aactgactca	cacttcccca	gacaggacca	gttgccctca	tttccaaaga	actgcactct	1080
ggaattaaag	ggactcttcc	attttgaaga	aggcatccag	aagctgtatc	agtgcaatgg	1140
gatcgcttgg	aaagcctgga	gtccccaac	caaggatgtg	gaagacaaat	cctgtccagc	1200
cgggtggcac	cagcactcag	gctactgtca	catcttgatc	acagagcaga	aaggcacctg	1260
gaatgcggct	gcccagctt	gcagggaaca	atacctgggc	aaccttgtaa	ctgtattctc	1320
caggcagcac	atgcggtggc	tctgggacat	tggtgggaga	aagtcctttt	ggataggttt	1380
gaagaccaca	gtgcattgctg	gccactggga	gtggatcggg	ggtgaacctg	ttgccttcac	1440
caatgggaga	agagggccct	ctccacgtc	caagcttggg	aagagctgtg	ttttggttca	1500
aagacaaggg	aaatggcaaa	caaaagactg	taggagagcc	aaacctcata	attatgtgtg	1560
ttccagaaaa	ctctaaatat	aacagaccct	acagggggcc	acctggagtt	tgacacctat	1620
ttatttcacag	gatctgtgaa	tattgtctca	tagaaaaaaa	attgttatga	ttgagtgggt	1680
atacctttgt	gattctgtct	agtgaatatg	ggacattttt	aatagtcca	gaaagattga	1740
taaataaata	ttttttacaa	gataagatac	aatttttgta	tctcaatacc	ttttaaaata	1800
aatgccagca	gtattaaaaa	aaaaa				1825

<210> 50
 <211> 1712
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4675668CB1

<400> 50
 cttttcttcag tccccacgtg cgatccttcc cggcaacttt ttcgagaaaa atgcccacaaat 60
 tcaaggcggc ccgtgggggtg ggggggtcagg aaaaacatgc gcccttgccc gatcagatcc 120
 tggctgggaa tgcgggtgcgg gcgggggtccc gggagaagcg gcgggggtcgc gggacaggag 180
 aagcggagga agagtatgtg gggccccggc tgagccgacg gattttgcag caagcacggc 240
 agcaacagga ggaactcgag gccgagcatg ggactgggga caagcccgcg gcgccgcggg 300
 aacgcaccac gcggctgggt ccaagaatgc ctcaggatgg atcagatgac gaggacgagg 360
 agtggccccc cctgggagaag gctgccacaa tgacagcagc gggccatcat gcagagggtg 420
 ttgtggaccc tgaggatgag cgtgccatag agatgttcat gaacaagaac cctcctgcca 480
 ggcgccacct ggctgacatc atcatggaga agctgactga gaagcagaca gaggttgaga 540
 cagtcatgtc agagggtgtc ggcttcccta tgccccagct ggaccccccg gtccctagaag 600
 tgtacagggg ggctcgggag gtattatcta agtaccgcag tggaaaactg cccaaggcat 660
 ttaagatcat ccctgcactc tccaactggg agcaaatcct ctacgtcaca gagccggagg 720
 cctggactgc agctgccatg taccaggcca ccaggatatt tgctctaac ctgaaggaaac 780
 gcatggccca gcgcttctac aaccttgtcc tgctccctcg agtacgagat gacgttgctg 840
 aatacaaagc actcaacttc catctctaca tggctctcaa gaaggccctt ttcaaacctg 900
 gagcctgggt caaagggatc ctgattccac tgtgcgagtc tggcacttgt accctcggg 960
 aagccatcat tgtgggtagc atcatcacca agtgctccat ccctgtgttg cactccagt 1020
 cggccatgct gaaaattgct gagatggaat acagcgggtg caacagcatc ttcttgcgac 1080
 tgctgctgga taagaagtat gcactgcctt accgggtgct ggatgcccta gtcttccact 1140
 tcctgggggt ccggacagag aagcgtgaac tgctgtgct gtggcaccag tgcctcctga 1200
 ctttgggtcca gcgtacaag gccgacttgg ccacagacca gaaagaggcc ctcttagaac 1260
 tgctccggct gcagcccat ccacagctat cggccgaaat caggcgtgag cttcagagt 1320
 cagtcccccg cgatgtggaa gatgttccca tcaccgtgga gtgaggaaaa cagtcagctg 1380
 tcctggccaa aggggttttg aaggacacca agaccccggt tgggtgactga agatgacact 1440
 gagctttaat ggctgaagac ccagatcagg gcagtgcag atcacaggga catctgtggc 1500
 tcccagttca ggacaggaag gactgagggt ctggctgggt ccctcttcca ttctaggccc 1560
 ttatccctgt ttagttctga gagccaact gagataccat atgctagcat tcccagttcc 1620
 cagctggggc ttggtgtgag tactttttct atggctattg tgtcaggtca ctgtggataa 1680
 aggcaaagac agatatttat tgaaaaaaaa aa 1712